



Contents lists available at ScienceDirect

Developmental and Comparative Immunology

journal homepage: www.elsevier.com/locate/dci

Comparative transcriptomics in three ladybird species supports a role for immunity in invasion biology

Heiko Vogel ^a, Henrike Schmidtberg ^b, Andreas Vilcinskis ^{b, c, *}

^a Max-Planck Institute for Chemical Ecology, Entomology Department, Hans-Knoell-Strasse 8, 07745 Jena, Germany

^b Justus-Liebig-University of Giessen, Institute for Insect Biotechnology, Heinrich-Buff-Ring 26-32, 35392 Giessen, Germany

^c Fraunhofer Institute for Molecular Biology and Applied Ecology, Department of Bioresources, Winchester Str. 2, 35395 Giessen, Germany

ARTICLE INFO

Article history:

Received 10 August 2016

Received in revised form

25 September 2016

Accepted 26 September 2016

Available online xxx

Keywords:

Invasion biology

Immunity

Transcriptomics

Ecoimmunology

Adalia bipunctata

Coccinella septempunctata

Harmonia axyridis

ABSTRACT

The spread of the invasive harlequin ladybird (*Harmonia axyridis*) in Europe is accompanied by the decline of the native and non-invasive two-spotted ladybird (*Adalia bipunctata*). Here we show that microsporidia carried by *H. axyridis* can kill *A. bipunctata* following the oral uptake of spores, suggesting that their horizontal transmission via intraguild predation may help the invader to outcompete its native competitor. The native seven-spotted ladybird (*Coccinella septempunctata*) is thought to be less susceptible both to the spread of *H. axyridis* and to its microsporidia. To investigate whether the distinct levels of pathogen susceptibility in these three ladybird species are determined by their immune systems, we compared the immunity-related transcriptomes of untreated beetles and beetles challenged with suspensions of bacteria and yeast. We found that *H. axyridis* carries three and four times as many genes encoding antimicrobial peptides representing the attacin, coleopteracin and defensin families than *C. septempunctata* and *A. bipunctata*, respectively. Gene expression studies following the injection of bacteria and yeasts into beetles revealed that members of these three antimicrobial peptide families are also induced more strongly in *H. axyridis* than *C. septempunctata* or *A. bipunctata*. Our results therefore support the hypothesis that a superior immune system promotes the performance of invasive species.

© 2016 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Invasive species represent a growing threat for biodiversity and the economy, but it remains unclear why some species are successful invaders whereas others, even closely related ones, are not (Tayeh et al., 2015). The harlequin ladybird (*Harmonia axyridis*) is recognized as a powerful model to evaluate factors that promote invasive success (Roy and Wajnberg, 2008). The native range of *H. axyridis* is central Asia, but it has been introduced into Northern America and Europe as an efficient biological control agent (Lombaert et al., 2011, 2014). It has become an invasive species in these new locations, and its spread is associated with the decline of native competitors such as the two-spotted ladybird (*Adalia bipunctata*) in the United Kingdom (Roy and Brown, 2015). Both *A. bipunctata* and the seven-spotted ladybird (*Coccinella septempunctata*) are predators that compete with *H. axyridis* for food in its

newly-colonized habitats (Pell et al., 2008; Kajita et al., 2010). All three species feed on each other's eggs and larvae, a phenomenon known as intraguild predation, which facilitates the horizontal transmission of pathogens among different ladybird species (Saito and Bjørnson, 2006; Vilcinskis, 2015).

Parasites co-introduced with alien species have been postulated to play a role in the outcome of biological invasions (Blackburn and Ewen, 2016). *H. axyridis* carries abundant spores of obligate parasitic microsporidia that do not harm the invasive vector but can kill the native ladybird *C. septempunctata* when injected (Vilcinskis et al., 2013a). To test our hypothesis that the microsporidia carried by *H. axyridis* function like bioweapons against native competitors, and can be transmitted via intraguild predation, we fed *A. bipunctata* larvae with *H. axyridis* eggs, larvae or isolated microsporidia, the latter added to the standard diet comprising eggs of the angoumois grain moth, *Sitotroga cerealella* (Vilcinskis, 2015). The mortality observed in the *A. bipunctata* population following the consumption of *H. axyridis* microsporidia suggests susceptibility to these pathogens in all three ladybird species is determined by their immune systems.

Ecoimmunology has emerged as a novel field which addresses

* Corresponding author. Institute for Insect Biotechnology, Justus-Liebig-University of Giessen, Heinrich-Buff-Ring 26-32, D-35392 Giessen, Germany.

E-mail address: andreas.vilcinskis@agr.uni-giessen.de (A. Vilcinskis).

<http://dx.doi.org/10.1016/j.dci.2016.09.015>

0145-305X/© 2016 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

also the role of immunity in invasion biology (White and Perkins, 2012). The immune systems of invasive species are thought to be superior to those of closely related non-invasive species (Lee and Klasing, 2004). This hypothesis is supported by the observation that *H. axyridis* is less susceptible to the entomopathogenic fungus *Beauveria bassiana* than the native ladybird species *A. bipunctata* and *C. septempunctata* (Roy et al., 2008). The robust antimicrobial defense of *H. axyridis* relies on both the constitutively present alkaloid (17R,9Z)-1,17-diaminooctadec-9-ene (harmonine) and antimicrobial peptides (AMPs) that are inducible by pathogens (Schmidtberg et al., 2013). Harmonine displays broad-spectrum activity against pathogens including human parasites such as *Leishmania major* and *Plasmodium falciparum* (Röhrich et al., 2012; Nagel et al., 2015). Next-generation-sequencing of the immunity-related transcriptome of *H. axyridis* led to the discovery of more than 50 genes encoding putative AMPs (Vilcinskas et al., 2013b). In order to determine whether this remarkable expansion in the repertoire of immunity-related effector genes evolved within the Coccinellidae, we compared the transcriptomes of untreated *A. bipunctata* or *C. septempunctata* beetles and beetles challenged with microbial pathogens. The resulting datasets were screened for the presence of AMPs and their expression levels before and after microbial challenge were compared to our previous comparative analysis of naïve and challenged *H. axyridis* beetles (Vilcinskas et al., 2013b).

2. Materials and methods

2.1. Insect material and RNA isolation

H. axyridis and *C. septempunctata* adults were collected in and around Giessen, Jena and Ober-Moerlen, Germany, for captive breeding. *A. bipunctata* adults were purchased from BIOCARE GmbH (Urbach, Germany). The rearing of all ladybird species, sample preparation, RNA isolation and the injection of suspensions containing 10 mg/ml lyophilized *Escherichia coli*, *Micrococcus luteus* and *Saccharomyces cerevisiae* were carried out as previously described (Vilcinskas et al., 2013a,b). RNA integrity was determined using an Agilent 2100 Bioanalyzer and RNA Nanochip (Agilent Technologies, Palo Alto, CA, USA), and RNA quantity was determined using a Nanodrop ND-1000 spectrophotometer.

2.2. Infection with microsporidia

A. bipunctata first to second larval instars were isolated in small Petri dishes and assigned to four groups with different diets: I) untreated *S. cerealella* eggs as a control diet; II) fresh *H. axyridis* eggs; III) *S. cerealella* eggs spiked with microsporidia isolated from *H. axyridis* (17.5×10^8 spores/65 mg eggs); and IV) living *H. axyridis* larvae. For the third diet, hemolymph was isolated from adult *H. axyridis* beetles by cutting the legs at the coxal base and drawing the hemolymph into two volumes of ice-cold anti-coagulant saline. The samples were then centrifuged at $1500 \times g$ for 10 min at 4 °C and the pellets were resuspended in phosphate buffered saline (PBS). The centrifugation and resuspension steps were carried out four times, and 5 µl of the final suspension was added dropwise to 50–100 *S. cerealella* eggs for each *A. bipunctata* larva (Vilcinskas et al., 2013a). During the tests, the different diets were renewed every second day. The survival rates were calculated using Kaplan-Meier survival analysis and the log-rank test in SigmaPlot v11 (Systat Software, Inc., San Jose, CA, USA). Microscopic inspection of dead *A. bipunctata* was performed to confirm the presence of microsporidia.

2.3. Transcriptome sequencing, assembly and annotation

A. bipunctata and *C. septempunctata* beetles were challenged by injecting a suspension containing 10 mg/ml lyophilized *E. coli*, *M. luteus* and *S. cerevisiae* as previously described (Vilcinskas et al., 2013a,b). Twenty four hours upon injection of microbial elicitors total RNA was isolated from 10 pooled beetles of each species in the naïve and challenged groups. Transcriptome sequencing was carried out by the Max Planck Genome Center Cologne (MPGCC) on an Illumina HiSeq2500 Genome Analyzer platform using poly(A)⁺ enriched RNA fragmented to an average of 150 nucleotides. This yielded approximately 30 million paired-end (2×100 bp) reads for each of the four samples. Quality control measures, including the filtering of high-quality reads and trimming the read length, as well as *de novo* transcriptome assemblies, were carried out using CLC Genomics Workbench v7.1 (<http://www.clcbio.com>), by selecting the presumed optimal consensus transcriptome and using previously described parameters (Vogel et al., 2014; Jacobs et al., 2016). The transcriptomes were annotated using BLAST, Gene Ontology and InterProScan searches implemented in BLAST2GO PRO v2.6.1 (www.blast2go.de) as previously described (Vogel et al., 2014). To identify *C. septempunctata* and *A. bipunctata* candidate AMP and lysozyme genes, we established a reference set of known or predicted insect-derived AMPs and lysozymes using published sequences and by searching our in-house database as well as public databases (NCBI). To avoid interpreting incomplete genes or allelic variants as different AMP genes, we used a number of additional filters to obtain a non-redundant set of candidate AMP genes using previously described criteria (Jacobs et al., 2016).

2.4. Sequence submission

We have deposited the short read (Illumina HiSeq2500) data to the European Nucleotide Archive (ENA) with the following study accession numbers: *A. bipunctata*: PRJEB11874 and *C. septempunctata*: PRJEB11877 (EBI short read archive (SRA)). The complete study is also accessible directly at the following URLs: <http://www.ebi.ac.uk/ena/data/view/PRJEB11874> and <http://www.ebi.ac.uk/ena/data/view/PRJEB11877>.

2.5. Analysis of AMP gene expression

Digital gene expression analysis was carried out using CLC Genomics Workbench v7.1 to generate BAM mapping files, QSeq (DNASTar, Inc., Madison, WI, USA) to remap the Illumina reads of each sample onto the corresponding reference transcriptomes, and finally by counting the sequences to estimate the expression levels, using previously described parameters for read mapping and normalization (Vogel et al., 2014; Jacobs et al., 2016). Briefly, the following parameters were used for read mapping: read assignment quality options required at least 50% of the total read bases and at least 90% of bases matching within each read to be assigned to a specific contig; maximum number of hits for a read = 10; mer repeat settings were automatically determined while other settings were not changed.

3. Results

3.1. Infection with microsporidia

A. bipunctata larvae were fed on the four diets described above and observed daily for mortality. The groups fed on *H. axyridis* eggs or first instar larvae died off rapidly (most individuals were dead within 2 weeks) and we determined microsporidia in the cadavers whereas the control group fed on *S. cerealella* eggs survived much

Download English Version:

<https://daneshyari.com/en/article/5540248>

Download Persian Version:

<https://daneshyari.com/article/5540248>

[Daneshyari.com](https://daneshyari.com)