



Short communication

Pharmacokinetics of a novel spot-on formulation of praziquantel for dogs



María Laura Gutiérrez^{a,b,c}, Guillermo Di Federico^d, Jorge A. Dale^e, Juan M. Minoia^c,
Carlos D. Corrales^e, Paula Schaiquevich^{a,f}, Silvia Wikinski^{a,b,c,*}

^a CONICET, Argentina

^b Instituto de Farmacología, Facultad de Medicina, Universidad de Buenos Aires, Paraguay 2155, C1121ABF Ciudad Autónoma de Buenos Aires, Argentina

^c Plataforma Tecnológica EBAL, Paraguay 2155, C1121ABF Ciudad Autónoma de Buenos Aires, Argentina

^d Servicio de Cirugía, Hospital Escuela, Facultad de Ciencias Veterinarias, Universidad Nacional de La Plata, Calle 60 y 118, 1900 La Plata, Provincia de Buenos Aires, Argentina

^e Laboratorio Labyes, Intendente Abel Costa 833, B1708JIN Morón, Provincia de Buenos Aires, Argentina

^f Unidad de Farmacocinética Clínica, Hospital de Pediatría J.P. Garrahan, Combate de los Pozos 1800, C1245AAL Ciudad Autónoma de Buenos Aires, Argentina

ARTICLE INFO

Keywords:

Anthelmintic
Transdermal route
Bioavailability
Dog

ABSTRACT

Praziquantel (PZQ) is an anthelmintic drug used both in humans and animals that can be administered through various routes. There are transdermal formulations for cats, but only oral or subcutaneous dosage forms for dogs. Given the fact that the cat's skin and the dog's skin have different characteristics, which in turn affect bioavailability, we developed a PZQ spot-on formulation for dogs. This study was aimed at determining the plasmatic behavior of topically administered PZQ (Labyes[®]) in adult dogs.

Dogs were administered PZQ (14.5 mg/kg PZQ, from a solution of 100 mg/ml). Blood samples were drawn before treatment onset and at the following time points after PZQ administration: 1, 2, 4, 6, 12, 24 and 48 h. PZQ plasma concentration was determined by ultra-high performance liquid chromatography (UPLC) coupled to tandem mass spectrometry (MS/MS). Observed maximum concentration (C_{max}), area under the concentration–time curve from the time of drug administration to infinity (AUC_{inf}) and time to maximum concentration (T_{max}) were calculated for each animal, and mean \pm SD for each parameter was obtained. Results were as follows: $C_{max} = 56.0 \pm 15$ ng/ml; $AUC_{inf} = 910.2 \pm 220$ ng* h/ml, $T_{max} = 5.0 \pm 1.1$ h. This is the first study to provide pharmacokinetic data of a praziquantel spot-on formulation for dogs.

1. Introduction

Praziquantel (PZQ), a synthetic isoquinoline-pyrazine derivative, is an anthelmintic used for the treatment of infections caused by various species of trematodes and cestodes in domestic animals. There are available both oral and subcutaneous dosage forms for administration of PZQ to dogs, but not transdermal formulations, which do exist for administration to cats.

The transdermal route of administration achieved via spot-on formulations has several advantages over the oral or subcutaneous forms; among them are the possibility of a controlled administration, the minimization of hazardous animal handling by the operator and a better tolerability for the animal, as it reduces the risk of vomiting and of pill rejection. Furthermore, the simplicity of the administration procedure shall very likely contribute to treatment adherence, thus increasing treatment effectivity. Regarding this matter, a critical analysis of the effectiveness of cystic echinococcosis eradication programs in South America demonstrated that the use of PZQ for 30

years has not been able to eradicate the endemic disease in dogs, nor its transmission to humans, mainly due to the practical complications of a large-scale and periodic treatment (Larrieu and Zanini, 2012).

From a pharmacokinetic standpoint, the transdermal route has the advantage of avoiding the first-pass effect, thus increasing drug bioavailability. It is also an adequate route for antiparasitic drug combinations, a formulation strategy that increases the chance of a thorough deworming in a single application.

The efficacy of transdermal PZQ, either alone or in combination with other endectocides, has been widely demonstrated in cats (Böhm et al., 2015; Jenkins and Romig, 2000; Krüdewagen et al., 2015; Traversa et al., 2009). However, its penetration in dogs remains to be determined.

Species-specific formulations need to be developed, due to the differences between the skin barriers of dogs and cats. Variables such as blood flow, capillary density, hair coat structure and hair growth rate, number and distribution of sebaceous glands and mainly skin thickness (21.16 ± 2.55 μ m in dogs versus 12.97 ± 0.93 μ m in cats)

* Corresponding author at: Paraguay 2155, 9th floor, C1121ABF Ciudad Autónoma de Buenos Aires, Argentina.
E-mail address: silviawikinski@gmail.com (S. Wikinski).

considerably affect drug absorption (Monteiro-Riviere et al., 2008).

In view of the above, we decided to develop a topical formulation for dogs, with characteristics similar to those of the ones already available for cats. We here present the first stage of our development; a demonstration of the systemic bioavailability of spot-on PZQ administered to dogs.

2. Materials and methods

2.1. Ethics committee's approval

This clinical trial was approved by the Ethics Committee of the Technological Platform EBAL (University of Buenos Aires-CONICET-University of La Plata). Informed consent was signed by the caregivers of the animals. The study was conducted according to Good Clinical Practices and International Guiding Principles for Biomedical Research Involving Animals as issued by the Council for the International Organizations of Medical Sciences.

2.2. Selection and inclusion of experimental subjects

Seven mixed-breed healthy adult dogs (5 males and 2 females), from the S.A.D.A.C. Animal Shelter located in Quilmes, Province of Buenos Aires, Argentina, were included in this trial. Mean \pm SD body weight values were 18.03 ± 4.8 kg, ranging from 12.7 to 25.0 kg.

Animals were physically examined by a veterinarian and health status was determined through the following parameters: nutritional status (appropriate weight for breed, gender and age, with no signs of obesity or of muscle loss), mucous membrane color, capillary refill time, behavioral parameters (aggression, submission, lethargy, tremors), psychomotor performance (normal, inactive, slowed-down), gait (rigidity, ataxia), heart and lung auscultation, abdominal palpation, presence or absence of disease-related signs (fever, vomiting, stool character).

Pregnant or lactating animals were excluded, as were underweight subjects or animals with clinical evidence of any disease, especially parasitic diseases.

Animals were individually housed in 7 suitable stainless steel kennels. Food was available, as provided in the Animal Shelter, in two daily servings, one of them consisting of rice, chicken and beef with bone and the other one of standard dry food; tap water was available *ad libitum*.

Subjects were identified with numbered collars; kennels were also numbered.

A clinical record card was kept, which contained subject number, kennel number, gender, weight, age, hair (short, medium, long), distinctive features, study drug dosage and adverse events.

2.3. Treatment and sample collection

Six of the seven dogs included in the study received a 14.5 mg/kg spot-on dose of PZQ provided by Chemo Group in a solution of 100 mg/ml (0.145 mg/kg). Thus, the volume administered was 0.145 ml/kg. The formulation consisted of a 10% PZQ solution prepared in a diluent that contained several antioxidant substances and a specific enhancer (aprotic polar solvent) (patent pending in the National Institute of Industrial Property, Argentina, May 2016). Animal #7 received no treatment and was used as negative control.

PZQ was applied topically directly on the animal's dry skin, in the back cervical area of the neck, pulling the hair apart to avoid significant spillage on the hair coat. This procedure prevents dogs from licking the product from the administration site.

One-milliliter blood samples were drawn either from the cephalic or the femoral vein before and 1, 2, 4, 6, 12, 24 and 48 h after drug administration. Samples were collected in tubes containing dipotassium EDTA as anticoagulant. Within two hours of collection, samples were

transferred to the laboratory, where plasma was separated by centrifugation and stored at -20 °C until processed.

2.4. Plasma PZQ quantitation method

Quantitation of PZQ in plasma was performed at a laboratory accredited by the College of American Pathologists. Briefly, plasma samples were pretreated with acetonitrile at a low pH to precipitate proteins. The supernatant was analyzed by ultra high performance liquid chromatography (UPLC) coupled to tandem mass spectrometry (MS/MS) detection. Plasma PZQ identification and quantitation was done by associating the ion mass spectrum obtained with its chemical structure, using the precursor ion $(M H)^+$ (m/z : 313.1).

2.5. Method validation

In order to validate the plasma PZQ quantitation method, the following parameters were evaluated; acceptance criteria are indicated between brackets: specificity and carryover, recovery, accuracy (percent relative error of less than 15%), precision (coefficient of variation of less than 15%), quantitation limits ($\geq 3 \leq 1000$ ng/ml), linearity (three plots with a minimum of 6 points each and R^2 greater than 0.98), robustness, sample stability through freeze/thaw cycles, short term stability, analytical conditions, storage, autosampler use, stability of the standard and long term stability. All parameters passed the acceptance criteria.

2.6. Data analysis

A non compartmental analysis was performed to calculate individual C_{max} , AUC_{inf} and T_{max} , for each subject. Mean (SD; upper and lower CI 95%) for each parameter were then calculated from the individual values.

3. Results

Fig. 1A shows the individual plasma PZQ concentration versus time profiles obtained for each experimental subject. Also, mean (SD) values are represented in Fig. 1B.

Calculated mean values (SD; 95% CI) of praziquantel pharmacokinetic parameters were as follows: $C_{max} = 56.0$ ng/ml (15; 40.4–71.7), $AUC_{inf} = 910.2$ ng* h/ml (220; 679–1141), and $T_{max} = 5.0$ h (1.1; 3.8–6.1).

4. Discussion

We here present the first study to provide pharmacokinetic data of a praziquantel spot-on formulation for dogs. Our results show that the formulation has a T_{max} of 5 h, with a range between 4 and 6 h, and that there are detectable plasma concentrations after 48 h of the administration. The maximum plasma concentration was 56.0 ± 15 ng/ml, attained after administering a dose of 14.5 mg/kg. Mean value of the area under the concentration-time curve from timepoint 0 to infinity was 910.2 ± 220 ng h/ml.

PZQ has been formulated both as a single drug and in combination with other antiparasitic drugs in spot-on preparations for cats (Krüdwagen et al., 2015; Kvaternick et al., 2014; Tielemans et al., 2014). According to the Profender™ drug monograph submitted by Bayer Laboratories to the European Medicines Agency, this product contains 85.71 mg/ml PZQ, and the obtained C_{max} for a dose similar to the one used in this trial (0.145 ml/kg) was of 61.3 ± 44.1 ng/ml (EMA, 2008), a value comparable to the one obtained by us. On the other hand, the T_{max} reported for Profender™ was 18.7 ± 47 h, a much greater value than the one found in our trial.

PZQ can also be found in another spot-on antiparasitic formulation for cats, Broadline™ from Merial laboratories. The dose in this case is

Download English Version:

<https://daneshyari.com/en/article/5545785>

Download Persian Version:

<https://daneshyari.com/article/5545785>

[Daneshyari.com](https://daneshyari.com)