



Research paper

In vitro antiparasmodial activity of *Withania frutescens*—Solanaceae

Laila El Bouzidi^{a,*}, Widad Ben Bakrim^a, Valérie Mahiou^b, Nadine Azas^c, Mustapha Larhsini^a, Mohamed Markouk^a, Evelyne Ollivier^b, Khalid Bekkouche^a

^a Laboratory of Biotechnology, Protection and Valorization of Plant Resources, Phytochemistry and Pharmacology of Aromatic and Medicinal Plant Unit, Faculty of Science – Semlalia, Cadi Ayyad University, P.O. Box 2390, 40000, Marrakech, Morocco

^b Laboratory of Pharmacognosy and Ethnopharmacology, UMR-MD3, Faculty of Pharmacy, Aix-Marseille University, 27 Bd. Jean Moulin, CS 30064, 13385, Marseille Cedex 5, France

^c Laboratory of Parasitology, UMR MD3, Faculty of Pharmacy, Aix-Marseille University, 27 Bd. Jean Moulin, CS 30064, 13385, Marseille Cedex 5, France

ARTICLE INFO

Keywords:

Antiplasmodial activity

Withania frutescens

Plasmodium falciparum

Cytotoxicity

Selectivity index

ABSTRACT

Introduction: Drugs resistant *Plasmodium falciparum* is a recurring issue that threatens public health. New antiplasmodial drugs are needed to overcome this problem. The aim of this study was to characterize *in vitro* antiparasmodial activity of *Withania frutescens* (Solanaceae) against chloroquine-resistant *Plasmodium falciparum* strain.

Methods: The *in vitro* antiparasmodial activity of leaves and roots extracts from *Withania frutescens* was performed in 96 well plates and preliminary phytochemical analysis was performed for the active fractions. The toxicity of the plant extract against CHO (Chinese Hamster Ovary) cells was assessed using the tetrazolium salt MTT colorimetric assay. The selectivity index (SI) was calculated as the ratio between the cytotoxic and antiparasitic activities.

Results: The methanol extract of *W. frutescens* leaves showed a good antiparasmodial activity (CC₅₀ 18.1 µg/ml). Furthermore, ethyl acetate and butanol fractions showed promising *in vitro* antiparasmodial activity with a selectivity index of 3.1 and 2.4, respectively. The roots of *W. frutescens* were found to be inactive with a CC₅₀ value > 80 µg/ml.

Conclusion: The antiparasmodial activity of *W. frutescens* may in part be attributed to the presence of polyphenolic and flavonoid compounds. Based on our results, ethyl acetate and butanol leaves fractions could be considered as a promising source for the development of putative antiparasmodial drugs.

1. Introduction

Malaria caused by *Plasmodium* parasites is amongst many prevalent public health concerns in several tropical regions of the world. Amongst the five existing species of *Plasmodium* causing malaria in humans, *Plasmodium falciparum* is the most widely studied species since it causes a high number of deaths [1]. In spite of considerable control efforts, malaria remains a major cause of global morbidity and mortality in most tropical countries, especially those in sub-Saharan Africa [2,3]. Resistance to the majority of available antimalarial drugs has been reported in a growing number of countries worldwide and such resistance threatens future progress in malaria control [1]. Medicinal plants have been a reliable source of therapeutic agents and nowadays still represent an inexhaustible pool for the discovery of novel drug leads.

Withania frutescens is a perennial plant mainly distributed in Morocco, Algeria, Spain and Canary Islands [4]. Previous investigation on the pharmacology of this plant showed that its crude extracts

displayed an interesting bioactivity profile, possessing various bioactivities including a protective and curative action against carbon tetrachloride (CCl₄)-induced hepatotoxicity, neuroprotective properties, antioxidant and cytotoxic effects [5–8]. The major chemical constituents reported from *W. frutescens* are called withanolides. These compounds are structurally diverse steroidal compounds with an ergosterol skeleton in which C-22 and C-26 are oxidized to form a δ-lactone. Many of these withanolides are reported to be responsible for the wide array of pharmacological activities, including antitumor, cytotoxic, cancer preventive, antifeedant, anti-inflammatory, leishmanicidal and immune-modulating effects [7,9–13]. In our ongoing research on *W. frutescens*, we aim to explore its antiparasmodial potential in view to discover new drugs. This paper deals, for the first time, with the *in vitro* evaluation of the antiparasmodial activity of *W. frutescens* against a chloroquine-resistant strain of *Plasmodium falciparum*.

* Corresponding author.

E-mail address: laila_elbouzidi@yahoo.fr (L. El Bouzidi).

2. Materials and methods

2.1. Plant material

Withania frutescens (L.) Pauquy roots and leaves were collected in El Hassania (20 km from Marrakech), Morocco, in March 2009. The material was authenticated by Prof. A. Abbad, a plant taxonomist, in the Department of Biology, Faculty of Sciences Semailia, Cadi Ayyad University, Morocco. A voucher specimen (No. mnhm.bk.kh.1) has been deposited at the herbarium of Natural History Museum of Marrakech, Morocco.

2.2. Extracts and fractions preparation

The leaves and roots were separately air-dried in the shade at room temperature (28 °C) until no further changes in their weight was observed (10 days), at Cadi Ayyad University of Marrakech (Morocco), in the laboratory of biotechnology, protection and valorization of plant resources, phytochemistry and pharmacology of aromatic and medicinal unit, and then reduced to fine powder with a grinder. Thus, the powdered parts of plants (100 g each) were exhaustively extracted using a Soxhlet apparatus. The leaves were first defatted with *n*-hexane (400 ml). The roots and defatted leaves were Soxhlet extracted with MeOH for 72 h (400 ml). After removing the solvent under vacuum at 40 °C, the residues obtained from the leaves (25.7 g) and roots (26.3 g) were dissolved in distilled water, filtered and extracted successively with CH₂Cl₂, EtOAc and *n*-BuOH, then concentrated under vacuum to yield 2.7 g, 0.6 g and 3.4 g of the CH₂Cl₂, EtOAc and *n*-BuOH leaf fractions, respectively and 0.9 g, 0.15 g and 3.1 g of the CH₂Cl₂, EtOAc and *n*-BuOH root fractions, respectively.

2.3. Phytochemical analysis of the plant extracts using TLC

The major secondary metabolite classes such as tannins, alkaloids, flavonoids and steroidal lactones were analyzed in *W. frutescens* by thin layer chromatography (TLC), on silica gel 60 F254 Merck. The solvent system used was: hexane/CH₂Cl₂/MeOH (18/3/2, v/v/v) for hexane extract, CH₂Cl₂/MeOH (9/1, v/v) for dichloromethane fraction, EtOAc/MeOH/H₂O (100/20/10, v/v/v) for ethyl acetate fraction and CH₂Cl₂/MeOH/H₂O (65/35/5, v/v/v) for methanol extract and butanol fraction. Visualization of the TLC plates was achieved under UV at 254 and 365 nm and by spraying with NP/PEG reagent for flavonoids, ferric trichloride for tannins and Dragendorff's reagent for alkaloids [14]. The presence of steroidal lactones was revealed with 4-4 nitrobenzylpyridine (NBP) followed by ethylenediamine (EDA) [15].

2.4. Determination of total phenolic compounds

The total phenolic content (TPC) of leaf extracts and fractions was determined using the Folin-Ciocalteu assay [16]. Aliquots of 300 µl containing 100 µg of each sample, dissolved in MeOH, were introduced into test tubes, followed by 1.5 ml of a Folin-Ciocalteu's reagent (diluted 10 times) and 1.2 ml of sodium carbonate (7.5% w/v). The contents of the tubes were mixed and placed in dark for 30 min before the absorbance was measured at 765 nm. The analyses were performed in triplicate and the results were expressed, as *p*-coumaric acid equivalent (CAE), in mg per g of dry extract or fraction ($y = 0.065x + 0.057$; $r^2 = 0.984$).

2.5. Determination of total flavonoid content

The total flavonoid content of leaf extracts and fractions was determined by the aluminium chloride colorimetric method [17]. In brief, 200 µl of each concentration (5 and 2.5 mg/ml) of the samples in methanol were mixed with 800 µl of distilled water and 60 µl of aqueous solution of NaNO₂ (5%). The mixture was allowed to stand for 5 min

before the addition of 40 µl of the solution of AlCl₃, and the mixture was allowed to stand for 5 min. Then, 400 µl of 1 M of NaCO₂ and 500 µl of distilled water were added. The absorbance of all samples was measured at 510 nm. The total flavonoid content was calculated from a calibration curve, and the result was expressed as mg catechin equivalent per g dry weight.

2.6. In vitro antiplasmodial assay

In this study, *P. falciparum* strain, resistant to chloroquine was used in the *in vitro* culture. Cultures were maintained in fresh A+ human erythrocytes at 2.5% haematocrit in complete medium (RPMI 1640 with 25 mM HEPES, 25 mM NaHCO₃, 10% of A+ human serum) at 37 °C under reduced O₂ atmosphere (gas mixture 5% O₂, 5% CO₂, and 90% N₂). Parasitaemia was maintained daily between 1% and 6%.

The *P. falciparum* drug susceptibility test was carried out by comparing quantities of DNA in treated and control cultures of parasite in human erythrocytes according to a SYBR Green I fluorescence-based method using a 96-well fluorescence plate reader. Parasite culture was synchronized at ring stage with 5% sorbitol. Compounds were incubated in a total assay volume of 200 µl (RPMI, 2% haematocrit and 1% parasitaemia) for 72 h in a humidified atmosphere (5% O₂ and 5% CO₂) at 37 °C, in 96-well flat bottom plates.

Triplicate assays were performed for each sample. After incubation, the supernatant was discarded and cells were washed with 150 µl of PBS. 15 µl re-suspended cells were transferred to 96-well flat bottom non sterile black plates (Greiner Bio-one) already containing 15 µl of the SYBR Green lysis buffer (2X SYBR Green, 20 mM Tris base pH 7.5, 20 mM EDTA, 0.008% w/v saponin, 0.08% w/v Triton X-100). Negative control, treated by solvents (DMSO) and positive control (chloroquine) was added to each set of experiments. Plates were incubated for 15 min at 37 °C and then read on a TECAN Infinite F-200 spectrophotometer with excitation and emission wavelengths at 497 and 520 nm, respectively. The concentrations of compounds required to induce a 50% decrease of parasite growth (CC₅₀) were calculated from three independent experiments.

2.7. In vitro cytotoxicity evaluation on CHO cell line

The evaluation of the tested extracts cytotoxicity was conducted on the Chinese Hamster Ovary (CHO) cells according to the method described by Mosmann [18] with slight modifications. Briefly, cells in 100 µl of complete medium, [RPMI supplemented with 10% foetal bovine serum, 1% L-glutamine (200 mM) and 1% penicillin (100 U/ml)/streptomycin (100 µg/ml)], were inoculated into each well of 96-well plates and incubated at 37 °C in a humidified 6% CO₂, 14% O₂ and 80% N₂ atmosphere. After 24 h incubation, 100 µl of medium with various product concentrations was added and the plates were incubated for 72 h. At the end of the treatment and incubation, each plate-well was microscope-examined for detecting possible precipitate formation before the medium was aspirated from the wells. 10 µl of MTT solution (5 mg MTT/ml in PBS) were then added to each well with 100 µl of medium without foetal calf serum. Cells were incubated for 2 h at 37 °C to allow MTT oxidation by mitochondrial dehydrogenase in the viable cells. After this time, the MTT solution was removed and DMSO (100 µl) was added to dissolve the resulting blue formazan crystals. Plates were shaken vigorously (300 rpm) for 5 min. The absorbance was measured at 570 nm with a microplate spectrophotometer (ELX 808, IU, BIOTEK). DMSO was used as blank and doxorubicine as positive control. Cell viability was calculated as percentage of control (cells incubated without compound). The 50% cytotoxic concentration was determined from the dose-response curve.

2.8. Selectivity index (SI)

The selectivity index (SI), which corresponds to the ratio between

Download English Version:

<https://daneshyari.com/en/article/5547227>

Download Persian Version:

<https://daneshyari.com/article/5547227>

[Daneshyari.com](https://daneshyari.com)