



The distribution of the modified Rankin scale scores change according to eligibility criteria in acute ischemic stroke trials: A consideration for sample size calculations when using ordinal regression analysis



Verónica V. Olavarría^a, Alejandro Brunser^b, Norberto Cabral^c, Sheila Martins^d,
Paula Muñoz-Venturelli^{b,e}, Gabriel Cavada^f, Pablo M. Lavados^{g,*}

^a Unidad de Neurología Vascular, Servicio de Neurología, Departamento de Neurología y Psiquiatría y Departamento de Paciente Crítico, Clínica Alemana de Santiago, Facultad de Medicina Clínica Alemana Universidad del Desarrollo, Santiago, Chile

^b Unidad de Neurología Vascular, Servicio de Neurología, Departamento de Medicina Interna, Clínica Alemana de Santiago, Facultad de Medicina Clínica Alemana Universidad del Desarrollo, Santiago, Chile

^c Universidade de Joinville (Univille), Registro de AVC de Joinville, Clínica Neurológica de Joinville, Joinville, Brazil

^d Hospital de Clínicas de Porto Alegre, Porto Alegre, RS, Brazil

^e The George Institute for Global Health, The University of Sydney, Sydney, Australia

^f Unidad de Investigación y Ensayos Clínicos, Departamento Científico Docente, Clínica Alemana de Santiago, Santiago, Chile

^g Unidad de Neurología Vascular, Servicio de Neurología, Departamento de Neurología y Psiquiatría, Clínica Alemana de Santiago, Facultad de Medicina Clínica Alemana Universidad del Desarrollo, Santiago, Chile

ARTICLE INFO

Article history:

Received 4 August 2016

Received in revised form

19 December 2016

Accepted 10 January 2017

Available online 5 February 2017

Keywords:

Stroke

Sample size

Clinical trials

Ordinal logistic regression

ABSTRACT

Background: Distribution shifts of the modified Rankin scale (mRs) is used as outcome measure in acute ischemic stroke (AIS) randomized controlled trials (RCT). Distribution across strata of mRs is relevant for sample size calculations and may be affected by eligibility criteria.

Aim: We aimed to assess the distribution of mRs scores across its different strata in AIS according to usual eligibility criteria.

Methods: We computed follow-up mRs strata distribution between an unselected cohort and samples with (a) time from symptom onset < 6 h (b) National Institutes of Health Stroke Scale (NIHSS) scores > 3 and < 25, and (c) both criteria combined. We compared distributions with the Mann-Whitney *U* Test and calculated sample sizes for each distribution.

Results: We included 5849 AIS patients. The unselected sample had a non-normal distribution with a median of 2. All selection criteria yielded significantly different distributions of mRs ($p = 0.04, 0.02$ and 0.02 respectively). This resulted in a significant variation in the calculated sample size when applying different selection criteria, with smaller numbers when RCT selection criteria are used (3616 versus 1553).

Conclusions: The use of usual RCT eligibility criteria result in significant differences in mRs distribution and smaller sample sizes compared to unselected AIS samples.

© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author. Unidad de Neurología Vascular, Servicio de Neurología, Departamento de Neurología y Psiquiatría, Clínica Alemana de Santiago, Facultad de Medicina Clínica Alemana Universidad del Desarrollo y Departamento de Ciencias Neurológicas, Facultad de Medicina, Universidad de Chile, Santiago, Chile.

E-mail addresses: veroolavaria@yahoo.com (V.V. Olavarría), abrunser2002@yahoo.com (A. Brunser), norbertocabral@icloud.com (N. Cabral), smartins@portoweb.com.br (S. Martins), pmventurelli@gmail.com (P. Muñoz-Venturelli), gabrielcavada@gmail.com (G. Cavada), pablolavados@yahoo.com (P.M. Lavados).

1. Introduction

The modified Rankin scale (mRs) is an ordinal disability score of 7 categories (0 = no symptoms to 6 = dead) and is currently the preferred method of evaluating outcome in patients with acute ischemic stroke (AIS) after intervention trials, particularly if the effect size is modest and distributed across all strata [1–3]. Thus, sample size calculations are increasingly determined by the distribution of the scores in the mRs at 90 days after symptom onset in modern AIS randomized clinical trials (RCT) [4–6].

Table 1
Demographic and clinical characteristics of sample cohorts.

Variables	Clínica Alemana, Santiago N = 1078	Hospital das Clínicas, Porto Alegre N = 1830	Joinville N = 2941
Mean age (SD)	70 (16)	66 (14)	66 (14)
Female (%)	539 (50)	934 (51)	1375 (48)
Hypertension (%)	722 (67)	1492 (82)	2170 (74)
Diabetes Mellitus (%)	226 (21)	510 (28)	957 (33)
Hypercholesterolemia (%)	259 (24)	829 (45)	887 (30)
Current smoking (%)	205 (19)	373 (20)	602 (20)
Atrial fibrillation (%)	172 (16)	338 (19)	394 (10)
Ischemic heart disease (%)	197 (9)	418 (23)	170 (6)
Previous stroke/TIA (%)	259 (24)	653 (36)	988 (34)
Cardioembolic (%)	366 (34)	490 (27)	844 (29)
Atherothrombotic (%)	140 (13)	423 (23)	706 (24)
Lacunar (%)	119 (11)	377 (21)	625 (21)
Undetermined (%)	420 (39)	347 (19)	597 (20)
Other determined (%)	32 (3)	193 (10)	169 (6)
NIHSS Median (IQR)	5 (2–10)	6 (3–13)	4 (2–11)

The data are analyzed using the *t*-test for difference of means or the Mann-Whitney *U* test or its variants - the Robust Rank Test and the Ordinal Logistic Regression (OLR) [7,8]. The latter also known as Whitehead's Odds Ratio derives from the proportional odds model of McCullagh [9], and has been shown to produce the lowest sample sizes [10]. The analysis assumes an equal chance of being in any strata and requires that at least the proportion of subjects in each scale category in one of the groups be specified, leading to a same odds ratio independently of the chosen partitions [11]. Thus the distribution across strata is relevant in deciding which statistical method is most suitable for sample size calculations when analyzing mRs as an ordinal scale [12].

Acute ischemic stroke RCTs eligibility criteria may impact the outcome distribution across strata and have the potential of being significantly different when compared to unselected stroke patients and between different trials. This could affect the sample sizes to an unknown extent.

Our aim was to compare the distribution of follow-up mRs scores across its different strata in patients with acute ischemic stroke (AIS) from an unselected consecutive cohort with 3 groups of patients meeting usual RCTs eligibility criteria. We hypothesized that variations in the distribution of mRs would significantly influence the sample sizes, when using OLR.

2. Methods

In this retrospective cohort study, patients were selected from two prospective hospital stroke registries and one population-based registry (Joinville): two in Brazil (Porto Alegre and Joinville) and one in Chile (Santiago).

For the unselected cohort, we included all adult AIS patients (i.e. ≥ 18 years old) with clinical and imaging diagnosis of AIS, data available on NIHSS at admission, time from symptom onset to admission, mRs at follow up and informed consent given. Patients

with Transient Ischemic Attacks and with incomplete data were excluded.

The following eligibility criteria have been used in many acute stroke RCTs and were applied to the cohort to produce 3 samples: (a) time from symptom onset < 6 h (b) National Institutes of Health Stroke Scale (NIHSS) scores > 3 and < 25 , and (c) both criteria combined.

The corresponding institutional review board and ethics committee approved the 3 registries were these data were derived from: The RECCA registry in Clínica Alemana, the Joinville Stroke Registry and Hospital das Clínicas, Porto Alegre Stroke Registry.

2.1. Analysis

We computed skewness, kurtosis, median values and used Mann-Whitney *U* Test to compare mRs strata distribution for each set of criteria compared to the unselected sample. We then applied the method proposed by S. Simon to calculate sample sizes for an acute trial using the distribution of mRs according to the different sets of eligibility criteria [13]. We grouped categories mRs 5 and 6 into one, as is usual in the analysis of AIS RCTs. All sample sizes were calculated with a modest effect size of 2% between categories of mRs in each group, α was set at 0.05 and power at 80% and all calculation were 2 tailed.

3. Results

The total unselected cohort consisted of 5849 consecutive patients, the demographic and clinical characteristics of which are described for each individual sample in Table 1.

The selection criteria yielded 3275 (56.0%) patients for NIHSS alone, 2636 (45.1%) for time alone and 1134 (19.4%) for both criteria combined (Table 2).

The unselected sample had a non-normal distribution of mRs,

Table 2
Distribution of modified Rankin scores at follow up in the total cohort according to usual acute ischaemic stroke trials selection criteria.

mRs scores	Unselected cohort N (%)	NIHSS >3 and < 25 N (%)	Time from onset <6 h N (%)	Both criteria N (%)
0	1211 (20.1)	391 (11.9)	531 (20.1)	132 (11.6)
1	1664 (28.4)	694 (21.2)	656 (24.9)	258 (22.6)
2	695 (11.8)	447 (13.6)	283 (10.7)	136 (11.9)
3	620 (10.6)	486 (14.8)	278 (10.5)	142 (12.5)
4	628 (10.7)	539 (16.5)	318 (12.1)	167 (14.7)
5–6	1031 (17.6)	2718 (21.9)	570 (21.6)	299 (26.3)
Total	5849	3275	2636	1134

Download English Version:

<https://daneshyari.com/en/article/5549657>

Download Persian Version:

<https://daneshyari.com/article/5549657>

[Daneshyari.com](https://daneshyari.com)