Accepted Manuscript

Title: Development of nanostructured lipid carriers for intraoral delivery of Domperidone

Authors: Carolin Tetyczka, Martin Griesbacher, Markus Absenger-Novak, Eleonore Fröhlich, Eva Roblegg

PII: S0378-5173(17)30386-1

DOI: http://dx.doi.org/doi:10.1016/j.ijpharm.2017.04.076

Reference: IJP 16642

To appear in: International Journal of Pharmaceutics

Received date: 15-3-2017 Revised date: 25-4-2017 Accepted date: 28-4-2017

Please cite this article as: Tetyczka, Carolin, Griesbacher, Martin, Absenger-Novak, Markus, Fröhlich, Eleonore, Roblegg, Eva, Development of nanostructured lipid carriers for intraoral delivery of Domperidone.International Journal of Pharmaceutics http://dx.doi.org/10.1016/j.ijpharm.2017.04.076

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Development of nanostructured lipid carriers for intraoral delivery of Domperidone

Carolin Tetyczka^a, Martin Griesbacher^a, Markus Absenger-Novak^b, Eleonore Fröhlich^b, Eva Roblegg^{a,c*} eva.roblegg@uni-graz.at

^aUniversity of Graz, Institute of Pharmaceutical Sciences, Department of Pharmaceutical Technology and Biopharmacy, Graz, Austria


^bMedical University of Graz, Center for Medical Research, Graz, Austria

^cResearch Center Pharmaceutical Engineering, Graz, Austria

* Corresponding Author: University of Graz, Universitätsplatz 1, 8010 Graz, Austria.

Tel.: +43 316 380 8888; Fax: +43 316 380 9100

Graphical Abstract



Abstract

The oral cavity displays one potentially attractive route that is not associated with gastric transit and hepatic first pass metabolism. However, poorly soluble active candidates require a drug delivery system to facilitate their transport through oral biological membanes. To this end, nanostructured lipid carriers (NLC) were loaded with Domperidone and produced with high pressure homogenization. NLC were

Download English Version:

https://daneshyari.com/en/article/5550275

Download Persian Version:

https://daneshyari.com/article/5550275

<u>Daneshyari.com</u>