



Contents lists available at ScienceDirect

Saudi Pharmaceutical Journal

journal homepage: www.sciencedirect.com



Original article

## Impacts of triamcinolone acetonide on femoral head chondrocytic structures in lumbosacral plexus block



Dashou Wang<sup>a,\*</sup>, Qian Chen<sup>a</sup>, Fengjun Cai<sup>d</sup>, Qi Pan<sup>a</sup>, Xuesong Li<sup>c</sup>, Qianming Wu<sup>a</sup>, Yong Gan<sup>a</sup>, Fei Meng<sup>c</sup>, Ping Luo<sup>b</sup>

<sup>a</sup> Department of Pain Management, Guizhou Provincial Orthopaedic Hospital, Guiyang 550002, China

<sup>b</sup> Department of Critical Care Medicine, Guizhou Provincial Orthopaedic Hospital, Guiyang 550002, China

<sup>c</sup> Department of Pain Intervention, Affiliated 300 Hospital of Guizhou Aviation Group, AVIC, Guiyang 550007, China

<sup>d</sup> Department of Pain Intervention, The Third People's Hospital of Guizhou, Guiyang 500002, China

### ARTICLE INFO

#### Article history:

Available online 21 April 2017

#### Keywords:

Lumbosacral plexus block  
Triamcinolone acetonide  
Chondrocytes  
Optical microscope  
Nerve axonal motion

### ABSTRACT

**Objective:** To investigate impacts of triamcinolone acetonide (TRI) on femoral head chondrocytic (FHC) structures when used for lumbosacral plexus block (LPB). **Methods:** A total of 32 6-month-old New Zealand white rabbits were selected (averagely weighing 2.75–3.25 kg) and added TRI into nerve block solution for LPB. The rabbit were randomly divided into four groups: group A1: 2.5 ml × 2 times, group A2 2.5 ml × 4 times, group B1 5 ml × 2 times, and group B2 5 ml × 4 times; the time interval among the injection was 5 days, and the structural changes of FHC were the observed using 50/100/200 light microscope; the modified Mankin pathological scoring was also performed for the evaluation. **Results:** There exhibited significant microscopic changes of FHC structures between the rabbits performed LPB and the normal rabbits, among which group B2 exhibited the most serious FHC damages, and the Mankin pathological score in group B2 was much higher than those in the other three groups, and the scores of the experimental group were higher than the control group. **Conclusions:** The addition of TRI in LPB can damage the FHC structures, and large-dose (5 ml/once) and long-course (four times) will result in more serious injuries.

© 2017 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### 1. Introduction

Chronic pain always seriously affects patients' daily activities and quality of life (Dueñas et al., 2016), and the morbidity increases in recent years (Gordon and Bloxham, 2016; Abbas et al., 2017). Since prolonging the peripheral nerve block may reduce the pain (Liu et al., 2015a; YaDeau et al., 2008), nerve blockade is often performed as therapeutic or palliative interventions for pain relief, especially for chronic pain (Luo et al., 2016; Hayek and Shah, 2014).

Nowadays, interventional procedures of nerve block are a widely accepted modality for pain management in adults (Shah et al., 2016), which would reduce the adverse reactions

systemically (Wardhan, 2015; Ali et al., 2017). A meta-analysis compared nerve block with physical therapy and intra-articular injection which focused on shoulder pain, and revealed that among the 591 patients, nerve block is of priority than physical therapy and not less useless than intra-articular injection (Chang et al., 2016). Another study focused on coccygodynia and draw the conclusion that ganglion impair block appears to be more effective than conservative therapy in resistant patients, 82% of the patients achieve at least 50% relief of pain after the first treatment, and the relief lasts for a median duration of 6 months (Gunduz et al., 2015).

Nerve block used in the treatment of chronic pain has already been developed maturely, and achieves good effects for osteoarthritis of knee joint, hip osteochondritis, and early femoral head necrosis, and a single-injection of femoral nerve block has been recommended as a routine therapy (Cien et al., 2015), which is also suggested in mild neurological deficit (Jonayed et al., 2016).

At the same time, the value of glucocorticoid has been demonstrated, such as dexamethasone is helpful to prolong pain block duration (YaDeau et al., 2015). A meta-analysis found that the addition of dexamethasone into local anesthetics prolongs brachial

\* Corresponding author.

E-mail addresses: 460659001@qq.com, 020416wzj@163.com (D. Wang).

Peer review under responsibility of King Saud University.



plexus block (Choi et al., 2014). Even low-dose dexamethasone (1–2 mg) would prolong the analgesia duration and motor blockade (Yine et al., 2015; Zhao and Ashraf, 2016; Liu et al., 2015a,b; Ghafar et al., 2017).

The addition of triamcinolone acetonide (TRI) when performing nerve block can achieve very good effects in nerve block therapies. Nerves have internal axonal motions, and this motion can transport glucocorticoids into joints so as to play the roles of anti-inflammation, analgesia, and joint pain relief. However, it has not been reported whether the treatment of distal articular cartilages can cause damages. This study used one optical microscope to observe the articular cartilage, aiming to see if the articular cartilage was affected. The study is reported as follows.

## 2. Materials and methods

### 2.1. Animals

A total of 32 6-month-old New Zealand white rabbits, weighing 2.75–3.25 kg, with the average as 2.95 kg, were selected and then divided into four groups ( $n = 8$ , gender-unrestricted). This study was carried out in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. The animal use protocol has been reviewed and approved by the Institutional Animal Care and Use Committee (IACUC) of Guizhou Provincial Orthopaedic Hospital.

### 2.2. Experimental methods preliminary experiment

Three rabbits were firstly shaved the right waist fur, and then placed in the left lying position; the needling site was positioned 1.5 cm laterally away from the right side of the spinous process of spine-iliac connection, and then one needle punctured 2–2.5 cm so as to break through the intertransverse ligament and enter the intergroove of right greater psoas muscle. After confirmed the position accuracy using X-ray, 1 ml of iohexol was then injected, and then computed tomography (CT) was performed to check the distribution of the contrast agent within the intergroove of right greater psoas muscle, and the depth was also recorded. Another three rabbits were then selected and performed the positioning using the same method, with the depth referring to the average depth obtained in the first time experiment, during which period the sense of breaking through should also be reviewed; 2.5 ml and 5.0 ml of Methylthionium Chloride (MC) were then injected, respectively. The rabbits were then euthanized using propofol and dissected so as to inspect the location and distribution of MC. Once it was confirmed distributing within the intergroove of right greater psoas muscle, and the distribution of lumbosacral plexus block (LPB) can be seen, it can confirm the consistency of the two positioning methods (Fig. 1).

The rabbits of the two groups were placed in the left lying position, positioned, and then medicated into the intergroove of right greater psoas muscle; the muscle strength of lower extremities were then observed and classified according to the human muscular strength classification. The final concentrations for LPB were as follows: lidocaine 0.5%, VB12 50  $\mu\text{g/ml}$ , TRI 2 mg/ml. The doses and injection times of the four groups: group A1: 2.5 ml  $\times$  2 times, group A2 2.5 ml  $\times$  4 times, group B1 5 ml  $\times$  2 times, and group B2 5 ml  $\times$  4 times, with the injection interval is 5 days.

### 2.3. Detection indexes

One optical microscopy was used to observe the right femoral head chondrocytic (FHC), and the modified Mankin pathologic scoring was used to evaluate each group; group A1 and A2, B1



Fig. 1. The needle clearance of the body surface was determined by metal needle under the C arm.

and B2, A1 and B1, and A2 and B2 were performed statistical X2 analysis.

### 2.4. Statistical analysis

SPSS 16.0 software was used for statistical analysis. The counting data were analyzed using the  $t$  test; the measurement data were expressed as mean  $\pm$  standard deviation, using the chi-square test, with  $\alpha = 0.05$  and  $P < 0.05$  considered as a statistical difference and  $P < 0.01$  as a statistically significant difference.

## 3. Results

### 3.1. Determination of insertion point of puncture needle

The needling site was positioned 1.5 cm laterally away from the right side of the spinous process of spine-iliac connection, and then one needle punctured 2–2.5 cm so as to break through the intertransverse ligament and enter the intergroove of right greater psoas muscle. CT exhibited the contrast agent distributing paraspinally, and partially entering the retroperitoneum. After injecting 2.5 ml and 5 ml of MC, it can be seen that MC distributed up to the L2 level and down to the pelvic ilium. After treatment, the muscle strength of the two 2.5 ml groups was in class II–III, and that of the two 5 ml groups was in class II or so. All the rabbits can only walk with their legs dragging. No statistically significant difference can be seen among the groups under optical microscope.

### 3.2. Microscopic results

FHC tissue exhibited normal structures, smooth surface, uniform thickness, normal chondrocytic size, shape, number, and arrangement; no cartilage growth or ossification exhibited abnormality; the epiphyseal plate thickness was normal and uniform, the chondrocytic size, shape, number, and arrangement were normal, and no cartilage growth or ossification exhibited abnormality; the chondrocytic column was arranged regularly; the trabecular thickness and morphology were normal and closely connected; the osteoblasts and osteoclasts' size, shape, number, and arrangement were normal; the structure of bone marrow was normal, and the cell proportion and component composition showed no abnormality (Fig. 2).

Download English Version:

<https://daneshyari.com/en/article/5551485>

Download Persian Version:

<https://daneshyari.com/article/5551485>

[Daneshyari.com](https://daneshyari.com)