



Replication of the Zika virus in different iPSC-derived neuronal cells and implications to assess efficacy of antivirals



Kristina Lanko^a, Kristel Eggermont^b, Abdulsamie Patel^b, Suzanne Kaptein^a,
Leen Delang^a, Catherine M. Verfaillie^b, Johan Neyts^{a,*}

^a Laboratory of Virology and Experimental Chemotherapy, Rega Institute for Medical Research, University of Leuven (KU Leuven), Leuven 3000, Belgium

^b Stem Cell Institute, University of Leuven (KU Leuven), Leuven 3000, Belgium

ARTICLE INFO

Article history:

Received 27 March 2017

Received in revised form

29 June 2017

Accepted 13 July 2017

Available online 20 July 2017

Keywords:

Zika virus

iPSC

Antivirals

Neuronal cells

ABSTRACT

Infections with the Zika virus (ZIKV) are responsible for congenital abnormalities and neurological disorders. We here demonstrate that ZIKV productively infects three types of human iPSC (induced pluripotent stem cells)-derived cells from the neural lineage, i.e. cortical and motor neurons as well as astrocytes. ZIKV infection results in all three cell types in the production of infectious virus particles and induces cytopathic effects (CPE). In cortical and motor neurons, an Asian isolate (PRVABC59) produced roughly 10-fold more virus than the prototypic African strain (MR766 strain). Viral replication and CPE is efficiently inhibited by the nucleoside polymerase inhibitor 7-deaza-2'-C-methyladenosine (7DMA). However, ribavirin and favipiravir, two molecules that inhibit ZIKV replication in Vero cells, did not inhibit ZIKV replication in the neuronal cells. These results highlight the need to assess the potential antiviral activity of novel ZIKV inhibitors in stem cell derived neuronal cultures.

© 2017 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

The Zika virus (ZIKV) is a mosquito-borne flavivirus that was first isolated in 1947 in Uganda (Dick et al., 1952). The last years this previously neglected virus has spread through the Pacific islands to South America, where it is currently causing a large epidemic (WHO, 2016). Infection during pregnancy has been shown to result in an increased chance of fetal abnormalities such as microcephaly, brain calcifications and ophthalmological problems (de Paula Freitas et al., 2016; Mlakar et al., 2016; Oliveira Melo et al., 2016). In adults, neurological complications such as Guillain-Barré syndrome, meningoencephalitis and acute myelitis have been reported (Cao-Lormeau et al., 2016; Carteaux et al., 2016; Mécharles et al., 2016). According to the WHO, the ZIKV epidemic associated with microcephaly and neurological complications continues to represent a major challenge for public health, therefore the Strategic Response Plan was set up to prevent and manage the complications of ZIKV (WHO, 2016). The exact mechanism of vertical transmission, teratogenic effects and neuro-invasion remains to be explored. To date ZIKV has been detected in the amniotic fluid (RNA) (Calvet et al., 2016; Sarno et al., 2016) and fetal brain tissue

(RNA and viral particles) (Driggers et al., 2016; Mlakar et al., 2016) of fetuses that develop abnormalities upon ZIKV infection of the pregnant mothers. Several hypothesis of vertical transmission have been proposed (Adibi et al., 2016; Retallack et al., 2016; Tabata et al., 2016). Neural tropism is a characteristic of several flaviviruses: tick-borne encephalitis virus (TBEV) is among the common infective agents of meningoencephalitis in Europe, West Nile virus (WNV) and Japanese encephalitis virus (JEV) are known to cause encephalitis (Ludlow et al., 2016; Winkelmann et al., 2016), and neurological complications can also rarely occur in Dengue virus infection (Madi et al., 2014). In addition, TBEV has been found in brain tissue and spinal cord of infected patients (Gelpi et al., 2005), and WNV is associated with neuronal loss (Kelley et al., 2003). However, the birth defects caused by ZIKV have not been reported for other flavivirus infections.

To further explore biological processes underlying fetal infection and neurological complications, relevant cellular and animal models of ZIKV infection are being developed. Here we report on ZIKV infection, and inhibition thereof, in three types of induced pluripotent stem cell (iPSC)-derived neuronal cell types: cortical neurons, motor neurons and astrocytes. Cortical neurons form the human cerebral cortex responsible for higher nervous activity. The disturbance of proliferation and differentiation of these cells has been linked to various disorders including microcephaly (Manzini

* Corresponding author. Rega Institute for Medical Research, University of Leuven, Herestraat 49 box 1030, BE-3000 Leuven, Belgium.

E-mail address: johan.neyts@kuleuven.be (J. Neyts).

and Walsh, 2011). Together with cortical neurons, astrocytes represent the predominant cell population in the human brain. Among other functions, astrocytes are involved in the neuro-inflammatory response during central nervous system (CNS) infection (Hamel et al., 2017; Molofsky and Deneen, 2015). Motor neurons control effector organs such as the muscles. In the context of ZIKV infection numerous cases of motor disability have been reported (Cao-Lormeau et al., 2016; Mécharles et al., 2016).

To investigate which cell types might be primary targets of ZIKV infection in the nervous system, we differentiated wild-type human iPSC into early cortical neurons, motor neurons and astrocytes and studied the susceptibility of these different cell types for ZIKV infection. To this end, the prototypic ZIKV strain MR766 from the African lineage and the recent clinical strain PRVABC59 from the Asian lineage (Puerto Rico, December 2015) were used each at a multiplicity of infection (MOI) of $10E-4$ (as determined in Vero cells). We used a low inoculum since this may provide a more sensitive readout of susceptibility to infection than overloading cells with a high inoculum. Using a low inoculum also had the advantage that the special neuronal culture medium was not substantially diluted. Following infection, cell cultures were monitored microscopically and supernatant samples were collected at day 1, 3 and 6 post infection for cortical neurons and astrocytes and at day 1, 4 and 7 post infection for motor neurons until cytopathogenic effects (CPE) were observed. Viral RNA levels and the infectivity of viral progeny were quantified by means of qRT-PCR and end-point titration in Vero cells, respectively.

The three cell types proved susceptible to both ZIKV strains and produced infectious progeny virus (Fig. 1). The PRVABC59 strain produced significantly higher viral loads (both RNA and infectious virions) than the MR766 strain at the end of the experiment in

cortical neurons (day 6 p.i.) and in motor neurons (day 7 p.i.). No significant difference in viral loads between the two ZIKV strains was observed in astrocytes. The highest titers of infectious virus were observed in motor neurons (up to $10E+7$ TCID₅₀/ml for PRVABC59 and $10E+6$ TCID₅₀/ml for MR766); titers in cortical neurons increased up to $10E+6$ for PRVABC59 and $10E+5$ TCID₅₀/ml for MR766, whereas the lowest viral production was observed in astrocytes. Similar results were obtained by measuring viral genome copies in culture supernatants by qRT-PCR (Fig. 1). The dynamics of ZIKV replication was comparable in cortical neurons and astrocytes with a rapid increase of viral RNA load during the first 3 days and complete destruction of the cell cultures by CPE at day 6 p.i. The infection in motor neurons progressed slower with first signs of CPE appearing at day 5–6 p.i. and complete destruction of the cultures by CPE at day 7–8 p.i. Our findings are in line with recent other studies where ZIKV was shown to infect human neural progenitor cells (hNPCs) and early cortical neurons (Tang et al., 2016; Dang et al., 2016; Xu et al., 2016). Infection of astrocytes was also demonstrated (Xu et al., 2016; Qian et al., 2016). When using a 10-fold lower MOI ($10E-5$) cortical neurons and motor neurons still proved susceptible to ZIKV infection, whereas this MOI proved too low to result in a productive infection in astrocytes (as monitored over a period of 6 days) (Fig. S1).

We next studied the potential antiviral activity of 3 molecules for which we demonstrated earlier that they inhibit ZIKV replication in Vero cells (Zmurko et al., 2016) namely T-705 (favipiravir), 7-deaza-2'-C-methyladenosine (7DMA) and ribavirin (Table 1). T-705 is a broad-spectrum inhibitor with antiviral activity against many RNA viruses including flaviviruses [the West-Nile virus and the yellow fever virus (Furuta et al., 2013)], and 7DMA was initially developed as a polymerase inhibitor of the hepatitis C virus (Olsen




Fig. 1. hiPSC-derived cells of neural lineage are susceptible to ZIKV infection. Infectious virus and viral RNA levels of ZIKV MR766 (black) and PRVABC59 (grey) strains following infection of hiPSC-derived (A, D) cortical neurons, (B, E) motor neurons and (C, F) astrocytes as quantified by qRT-PCR or by end-point titration respectively. Data are mean values of three independent experiments and are presented as mean ($\pm$) SEM. * $p < 0.05$.

Download English Version:

<https://daneshyari.com/en/article/5551650>

Download Persian Version:

<https://daneshyari.com/article/5551650>

[Daneshyari.com](https://daneshyari.com)