Author's Accepted Manuscript

Flavonoid derivative (Fla-CN) inhibited adipocyte differentiation via activating AMPK and upregulating microRNA-27 in 3T3-L1 cells

Chun-Chun Gan, Tian-Wen Ni, Yang Yu, Nan Qin, Ying Chen, Mei-Na Jin, Hong-Quan Duan



 PII:
 S0014-2999(17)30006-7

 DOI:
 http://dx.doi.org/10.1016/j.ejphar.2017.01.009

 Reference:
 EJP71006

To appear in: European Journal of Pharmacology

Received date: 26 October 2016 Revised date: 3 January 2017 Accepted date: 11 January 2017

Cite this article as: Chun-Chun Gan, Tian-Wen Ni, Yang Yu, Nan Qin, Ying Chen, Mei-Na Jin and Hong-Quan Duan, Flavonoid derivative (Fla-CN) inhibite adipocyte differentiation via activating AMPK and up-regulating microRNA-2' in 3T3-L1 cells, *European Journal of Pharmacology* http://dx.doi.org/10.1016/j.ejphar.2017.01.009

This is a PDF file of an unedited manuscript that has been accepted fo publication. As a service to our customers we are providing this early version o the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

ACCEPTED MANUSCRIPT

Flavonoid derivative (Fla-CN) inhibited adipocyte differentiation via

activating AMPK and up-regulating microRNA-27 in 3T3-L1 cells

Chun-Chun Gan¹, Tian-Wen Ni¹, Yang Yu¹, Nan Qin^{1,2}, Ying Chen², Mei-Na Jin^{1*},

Hong-Quan Duan^{1,2*}

¹Tianjin Key Laboratory on Technologies Enabling Development Clinical

Therapeutics and Diagnostics (Theranostics), School of Pharmacy, Tianjin Medical

University, Tianjin 300070, People's Republic of China

²Research Center of Basic Medical Sciences, Tianjin Medical University, Tianjin

300070, People's Republic of China

jinmeina@tmu.edu.cn

duanhq@tmu.edu.cn

*correspondence. Tel.: +86 022 83336680, Fax: +86 022 83336560.

*correspondence. Tel.: +86 022 83336538, Fax: +86 022 83336538.

Abstract

Fla-CN (3-O-[(E)-4-(4-cyanophenyl)-2-oxobut-3-en-1-yl] kaempferol) is a semi-synthesized flavonoid derivative of tiliroside which exhibited anti-diabetic effect *in vivo*. Our previous study revealed the role of Fla-CN in anti-obesity and anti-diabetes *in vivo*, but the underlying mechanism remained to be addressed. The present study aimed to investigate the mechanism of anti-adipogenesis *in vitro*. Fla-CN markedly inhibited intracellular lipid accumulation in a dose-dependent manner, and the inhibitory effect was mainly limited to the early Download English Version:

https://daneshyari.com/en/article/5554839

Download Persian Version:

https://daneshyari.com/article/5554839

Daneshyari.com