Accepted Manuscript



High-fat diet aggravates 2,2′,4,4′-tetrabromodiphenyl ether-inhibited testosterone production via DAX-1 in Leydig cells in rats

Zhan Zhang, Yongquan Yu, Hengsen Xu, Chao Wang, Minghui Ji, Jun Gu, Lu Yang, Jiansheng Zhu, Huibin Dong, Shou-Lin Wang

PII: S0041-008X(17)30114-X
DOI: doi: 10.1016/j.taap.2017.03.010
Reference: YTAAP 13891

To appear in: *Toxicology and Applied Pharmacology*

Received date: 15 October 2016
Revised date: 2 March 2017
Accepted date: 10 March 2017

Please cite this article as: Zhan Zhang, Yongquan Yu, Hengsen Xu, Chao Wang, Minghui Ji, Jun Gu, Lu Yang, Jiansheng Zhu, Huibin Dong, Shou-Lin Wang , High-fat diet aggravates 2,2′,4,4′-tetrabromodiphenyl ether-inhibited testosterone production via DAX-1 in Leydig cells in rats. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Ytaap(2017), doi: 10.1016/j.taap.2017.03.010

High-fat diet aggravates 2,2',4,4'-tetrabromodiphenyl ether-inhibited testosterone production via DAX-1 in Leydig cells in rats

Zhan Zhang[1,2,4], Yongquan Yu[1,2,4], Hengsen Xu[2], Chao Wang[1,2], Minghui Ji[2], Jun Gu[2], Lu Yang[1,2], Jiansheng Zhu[1,2], Huibin Dong[3], Shou-Lin Wang[1,2,*]

[1] State Key Lab of Reproductive Medicine, Institute of Toxicology, Nanjing Medical University, 101 Longmian Avenue, Nanjing 211166, P. R. China

[2] Key Lab of Modern Toxicology of Ministry of Education, School of Public Health, Nanjing Medical University, 101 Longmian Avenue, Nanjing 211166, P. R. China

[3] Changzhou Center for Disease Control and Prevention, 203 Taishan Road, Changzhou, 2013022, P. R. China

[4] These authors contributed equally to this study

[*] To whom correspondence should be addressed: Key Lab of Modern Toxicology of Ministry of Education, School of Public Health, Nanjing Medical University, 101 Longmian Avenue, Nanjing 211166, P. R. China

Tel: +86-25-8686-8417

Fax: +86-25-8686-8499

E-mail: wangshl@njmu.edu.cn