


Accepted Manuscript

Title: Intoxication by gamma hydroxybutyrate and related analogues: clinical characteristics and comparison between pure intoxication and that combined with other substances of abuse

Authors: Òscar Miró, Miguel Galicia, Paul Dargan, Alison M. Dines, Isabelle Giraudon, Fridtjof Heyerdahl, Knut E. Hovda, Christopher Yates, David M. Wood



PII: S0378-4274(17)30216-3
DOI: <http://dx.doi.org/doi:10.1016/j.toxlet.2017.05.030>
Reference: TOXLET 9787

To appear in: *Toxicology Letters*

Received date: 14-2-2017
Revised date: 28-5-2017
Accepted date: 29-5-2017

Please cite this article as: Miró, Òscar, Galicia, Miguel, Dargan, Paul, Dines, Alison M., Giraudon, Isabelle, Heyerdahl, Fridtjof, Hovda, Knut E., Yates, Christopher, Wood, David M., Intoxication by gamma hydroxybutyrate and related analogues: clinical characteristics and comparison between pure intoxication and that combined with other substances of abuse. *Toxicology Letters* <http://dx.doi.org/10.1016/j.toxlet.2017.05.030>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Intoxication by gamma hydroxybutyrate and related analogues: clinical characteristics and comparison between pure intoxication and that combined with other substances of abuse

Òscar Miró^{1,*}, Miguel Galicia^{1,*}, Paul Dargan^{2,3}, Alison M. Dines², Isabelle Giraudon⁴, Fridtjof Heyerdahl⁵, Knut E. Hovda⁵, Christopher Yates⁶, Euro-DEN Research Group **, David M. Wood^{2,3}

¹Emergency Department, Hospital Clínic, IDIBAPS, University of Barcelona, Barcelona, Catalonia, Spain.

²Clinical Toxicology, Guy's and St Thomas' NHS Foundation Trust and King's Health Partners, London, UK.

³Clinical Toxicology, Faculty of Life Sciences and Medicine, King's College London, London, UK

⁴European Monitoring Centre for Drugs and Drug Addiction (EMCDDA), Lisbon, Portugal

⁵The National CBRNe Centre of Medicine, Department of Acute Medicine, Medical Division, Oslo University Hospital, Oslo, Norway.

⁶Emergency Department and Clinical Toxicology Unit, Hospital Universitari Son Espases, Mallorca, Spain.

*Miguel Galicia and Òscar Miró have equally contributed to this manuscript and should both be considered as first authors.

**Euro-DEN Research Group - Corporate Author: Evangelia Liakoni, Matthias E. Liechti, Gesche Jürgens, Carsten Boe Pedersen, Niall O'Connor, Gerard Markey, Adrian Moughty, Christopher Lee, Jacek Sein Anand, Jordi Puiguriquer, Catalina Homar, Florian Eyer, Odd Martin Vallersnes, Per Sverre Persett, Lucie Chevillard, Bruno Mégarbane, Raido Paasma, W Stephen Waring, Kristiina Pöld, Christian Rabe, Piotr Kabata.

Address for correspondence: Miguel Galicia, Emergency Department, Hospital Clínic, Barcelona, Catalonia, Spain. Telephone: (+34) 93.227.98.33. FAX: (+34) 93.227.56.93. Email: mgalicia@clinic.cat

Funding: The Euro-DEN Project was funded by the DPIP/ISEC Programme of the European Union; all of the authors had funding from the European Commission through the Euro-DEN project except ML and EL, whose costs were co-funded by the Swiss Centre of Applied Human Toxicology (SCAHT), KP and IG.

Highlights from "Intoxication by gamma hydroxybutyrate and related analogues: clinical characteristics and comparison between pure intoxication and that combined with other substances of abuse " (Manuscript ID TOXLET-D-17-00142).

- The 78.0% of patients were over 24 years of age (median 31 years) and 83.4% were men.
- The most co-used substances: ethanol, amphetamine and derivatives, cocaine, benzodiazepines.
- The most common symptoms: behaviour altered, agitation and reduced level of consciousness.
- Endotracheal intubation and mechanical ventilation was performed in 6.9% of the whole group.
- The GHB/GBL with co-intoxicants group have more severe symptoms and higher necessity of therapy.

Download English Version:

<https://daneshyari.com/en/article/5562172>

Download Persian Version:

<https://daneshyari.com/article/5562172>

[Daneshyari.com](https://daneshyari.com)