



RESEARCH PAPER

# Reliability of novice physiotherapists for measuring Cobb angle using a digital method



Patcharawan Suwannarat, PT, PhD<sup>a,b</sup>,  
Pattra Wattanapan, MD<sup>b,c</sup>, Arpassanan Wiyanad, PT, MSc<sup>a,b</sup>,  
Pakwipa Chokphukiao, PT, MSc<sup>a,b</sup>, Sininat Wilaichit, PT, MPH<sup>d</sup>,  
Sugalya Amatachaya, PT, PhD<sup>a,b,\*</sup>

<sup>a</sup> School of Physical Therapy, Faculty of Associated Medical Sciences, Khon Kaen University, Khon Kaen, Thailand

<sup>b</sup> Improvement of Physical Performance and Quality of Life (IPQ) Research Group, Khon Kaen University, Khon Kaen, Thailand

<sup>c</sup> School of Rehabilitation Medicine, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand

<sup>d</sup> Department of Physical Therapy, Damnoen Saduak Hospital, Ratchaburi, Thailand

## KEYWORDS

Cobb angle;  
spinal angle;  
reliability;  
spinal deformity;  
X-ray

**Abstract** *Background:* The Cobb's method is the most accurate and reliable method for kyphosis measurement. Conventionally, a sagittal Cobb angle was commonly derived from a lateral plain film. With computer technology, a digital method is widely used in common clinical settings, but the existing reliability data involved only experienced raters.

*Objectives:* To assess the interrater and intrarater reliability of a digital Cobb's method using novice physiotherapists.

*Methods:* Fifteen participants, with an occiput wall distance of more than 0 cm, were interviewed and assessed for their demographics. Then they were filmed for lateral spinal radiography over the area of thoracic spine in a standing position, and the Cobb angle was analyzed by four raters, including an expert physician and three novice physiotherapists, using a SurgimapSpine programme.

*Results:* The average Cobb angles among the four raters showed no significant difference ( $p = 0.984$ ). Outcomes of their measurements had excellent intrarater and interrater reliability [intraclass correlation coefficient ( $ICC_{3,3}$ ) = 0.995–0.997] with a small range of standard errors of the measurement ( $<1^\circ$ ).

*Conclusion:* A digital Cobb's method had excellent reliability when used by a novice health professional rater. The findings confirm the ease of using this method to detect and monitor kyphosis in general hospitals, clinics, or research facilities.

Copyright © 2017, Hong Kong Physiotherapy Association. Published by Elsevier (Singapore) Pte Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\* Corresponding author. School of Physical Therapy, Faculty of Associated Medical Sciences, Khon Kaen University, Khon Kaen, Thailand.  
E-mail address: [samata@kku.ac.th](mailto:samata@kku.ac.th) (S. Amatachaya).

## Introduction

The Cobb's method is one of the most common techniques to measure spinal curvature using radiograph. It was first described in 1948 by an American orthopedic surgeon, Dr John Robert Cobb, who outlined how to measure the angle of spinal curvature. Hence, the term "Cobb angle" came about bearing his name [1]. Originally, the Cobb angle has been used to measure coronal spinal deformity. Later, it was adapted to measure sagittal spinal angle, as the so-called kyphosis angle [2]. The method is accounted as the most accurate and reliable method [intra-class correlation coefficient (ICC) = 0.96–0.99]; thus, it is commonly used as a gold standard to validate a new kyphosis measure [3,4].

Previously, a sagittal Cobb angle was derived from a lateral plain film by drawing a straight line that passed the upper border of the fourth thoracic vertebra (T4), and another line that passed the inferior border of the 12<sup>th</sup> thoracic vertebra (T12). Then two other lines were drawn perpendicularly with the first two lines, and the angle of their intersection or the Cobb angle was measured using a protractor [4,5]. The intrarater and interrater reliability of this conventional Cobb's method has been reported among various raters including experienced physicians (ICC = 0.94), fellowships (ICC = 0.79–0.99), residents (ICC = 0.96–0.99), and rheumatologists (ICC = 0.91–0.95) [6–8]. However, with computer technology, a digital method offers a superior option for Cobb angle interpretation as compared to the conventional method. It allows data interpretation using a desktop computer, a laptop, or a smartphone, and reduces the cost for developing a plain film, storage areas for hard copies, and time required for data interpretation. In addition, digital software such as AutoCAD and SurgimapSpine offer a better view of the vertebra morphology through the adjustment of contrast and enlargement, which eases the measurement of distances and angles [9,10]. Consequently, various digital techniques are increasingly used in routine clinical assessments.

Currently, there is only a limited amount of reliability data of the digital Cobb's method, and all of them involve an experienced physician rater (ICC = 0.81–0.96) [9,11–13]. Apart from physicians, a physiotherapist is another important professional for kyphosis management to normalise or minimise the progression of kyphosis angle [5]. Briggs et al [14] have reported the reliability data of experienced physiotherapists who commonly evaluate Cobb's method. However, the reliability data of novice or inexperienced physiotherapists would confirm the ease of using a digital Cobb's method to detect and monitor kyphosis in general hospitals, clinics, or research institutions. Therefore, the current study aimed to evaluate the interrater and intrarater reliability of a digital Cobb's method that was measured by novice physiotherapist raters using computer-aided technique sagittal plane radiographs.

## Methods

### Participants

This study was cross-sectionally conducted from January to June 2016. The eligible participants were at least 18 years

old with an occiput wall distance (OWD) of >0 cm [15–17]. Exclusion criteria were any signs and symptoms of lower limb neurological compromises such as pain or inflammation in the muscles or joints, and other spinal or limb deformities that might confound data interpretation for spinal angles, i.e., scoliosis, amputation, and leg length discrepancy. The protocols of the study were approved by the Khon Kaen University Ethics Committee for Human Research, Khon Kaen, Thailand (HE581446).

### Research protocol

Walter et al [18] have suggested the method of sample size estimation for reliability studies using ICCs. The study set a true  $p_0$  at 0.4 (minimally acceptable level), an alternative  $p_1$  of 0.75, a 5% significance level, and a power of 80% ( $\beta = 0.20$ ). Thus, the study required at least 14 participants. The eligible participants were interviewed and assessed for their demographic characteristics, including age, sex, weight and height. Then, they were filmed for lateral spinal radiography over the area of the thoracic spine [the first thoracic vertebral (T1) to the 12<sup>th</sup> thoracic vertebral (T12)] in an upright standing position. The Cobb angle was subsequently analyzed by four raters using the SurgimapSpine software (version 1.2, Nemaris Inc, 306 East 15<sup>th</sup> St Suite 1R NY, New York 10003). The characteristics of the raters and details of the Cobb angle interpretation are presented in the following subsections.

### Characteristics of raters and training protocols for digital Cobb measurement

This study involved four raters, which comprised a physician who had extensive experience in digital Cobb's method (an expert) and three physiotherapists who did not have any experience in a digital Cobb measurement. At the beginning of the study, the three novice physiotherapist raters were trained by the expert. The training protocols consisted of didactic and demonstration in the areas relating to digital measure, including the basic spinal anatomy of the spine using data from an X-ray illustration, and instructions on (1) how to use the SurgimapSpine software, (2) how to find a landmark, and (3) how to read the angle from the programme. Then the three novice raters practiced using these methods until the expert was satisfied with their performance. In aggregate, the training time took approximately 45 minutes.

### Methods of Cobb angle measurement using a digital programme

To achieve a Cobb angle, a digital X-ray file was uploaded to the SurgimapSpine programme. Then, a straight line was drawn that passed the upper border of the T4 vertebra, and another line that passed the inferior border of the T12 vertebra. Then, two other lines were drawn perpendicularly with the first two lines, and the intersection of these two lines produced the Cobb angle [3,4]. Each rater carried out the measurements independently (three separate trials per rater), and the finding of each trial was blinded for a subsequent trial and for another rater.

Download English Version:

<https://daneshyari.com/en/article/5563838>

Download Persian Version:

<https://daneshyari.com/article/5563838>

[Daneshyari.com](https://daneshyari.com)