

Investigation of Preoperative Fasting Times in Children

Eda Dolgun, PhD, RN, Meryem Yavuz, PhD, RN, Birsen Eroğlu, RN, Ayşe Islamoğlu, RN

Purpose: *The aim of this study was to investigate preoperative fasting times of children.*

Design: *A prospective and descriptive design was used.*

Methods: *The study was carried out in the pediatric surgery departments of a University Faculty of Medicine Hospital, between January 1 and May 31, 2012. The sample included 332 pediatric patients having surgery with general anesthesia and their family. Data were collected via patient and family interview using a 28-item questionnaire developed by the researchers.*

Findings: *29.2% of the sample was between the age of 1 and 3 years, of which 68.1% were male. 82.5% had day surgery. 94.9% of families were informed about the operation and 97.6% about nutrition before the operation. The children were fed an average of 10.51 hours before the operation (n = 203) with clear liquids, 6.27 hours (n = 52) with breast milk, 9.9 hours (n = 107) with baby food or cow's milk, 11.22 hours (n = 100) with toast and other, and 12.25 hours (n = 106) with daily food. About 65.6% of children between the years 0 and 3 (n = 151) declared being hungry when they were going to surgery, and 58.9% said that they were thirsty. Of the children older than 3 years (n = 181), 55.2% were hungry and 58.0% were thirsty.*

Conclusions: *Results indicate that patients' preoperative fasting times were longer than recommended by current guidelines.*

Keywords: *fasting, preoperative care, pediatrics, perianesthesia nursing, preoperative period, research.*

© 2016 by American Society of PeriAnesthesia Nurses

Eda Dolgun, PhD, RN, is a Lecturer and Associate Professor at the Department of Surgical Nursing, Ege University Nursing Faculty, Bornova, İzmir, Türkiye; Meryem Yavuz, PhD, RN, is a Lecturer and Professor at the Department of Surgical Nursing, Ege University Nursing Faculty, Bornova, İzmir, Türkiye; Birsen Eroğlu, RN, is a Nurse at the Department of Pediatric Surgery, Ege University Faculty of Medicine, Bornova, İzmir, Türkiye; and Ayşe Islamoğlu, RN, is a Nurse at the Department of Pediatric Surgery, Ege University Faculty of Medicine, Bornova, İzmir, Türkiye.

Conflict of interest: None to report.

Address correspondence to Eda Dolgun, Department of Surgical Nursing, Ege University Nursing Faculty, Bornova, İzmir 35100, Türkiye; e-mail address: eda.dolgun@ege.edu.tr or edolgun@hotmail.com.

© 2016 by American Society of PeriAnesthesia Nurses

1089-9472/\$36.00

<http://dx.doi.org/10.1016/j.jopan.2014.12.005>

PERIANESTHESIA NURSES PLAY an important role in preparing children for surgery.¹ The preoperative fasting of children in the hospital is ensured because of nursing supervision. For outpatient surgical operations, however, preoperative fasting is the responsibility of the parents.²

The main aim of preoperative fasting is to reduce the risk of pulmonary aspiration of the stomach contents.¹⁻⁵ Prolonged preoperative fasting, however, might cause negative effects, such as a sense of hunger, discomfort, headache, dehydration, hypovolemia, and hypoglycemia.^{1,2,6} Staying hungry for a long time before an operation might also result in a decrease in patient satisfaction, late recovery, and late hospital discharge in the postoperative period.^{1,7}

Preoperative fasting guidelines recommend clear liquids up to 2 hours before surgery, breast feeding up to 4 hours, light meals up to 6 hours, and heavy/fatty meals for up to 8 hours before surgery. The suggestions differentiate depending on the age of the patient and the type of the food eaten.⁸⁻¹⁵

Although safe preoperative fasting times are defined, the issue remained unresolved in the study clinics. This study was designed to investigate preoperative fasting times and the hunger/thirst state of children undergoing general anesthesia.

Materials and Methods

Sample

This prospective descriptive study was carried out in the Pediatric Surgery Departments of a University Faculty of Medicine Hospital between January 1 and May 31, 2012. The sample included 332 patients who had surgery under general anesthesia and agreed to participate in the study.

Data Collection Tools

The data were collected using a 28-item questionnaire developed by the researchers. The questionnaire included nine questions about sociodemographic factors and 19 questions about pediatric operations, preoperative information received, nutritional conditions, and preoperative fasting time. Data collection was conducted via patient and family interviews. The level of hunger and thirst was ranked using a 0-to-10-point scale by children who were older than 3 years and by the parents of children between 0 and 3 years. Data were collected before the patients left the clinic. The time of assessment took approximately 15 minutes.

Ethical Aspects

The verbal consent of the families participating in the study and written consent of the University Faculty of Medicine Hospital was obtained to perform the study. Data were collected from the parents and children who were willing to participate in the study.

Statistical Analysis

Data analysis was performed using SPSS, version 16 (SPSS Inc., Chicago, IL), for Windows.

Findings

Sociodemographic factors are presented in [Table 1](#). 82.5% of the children had outpatient surgeries; 94.6% of parents were informed about the operation and dietary changes that would apply before the surgery.

The average preoperative solid meal fasting time was 10.17 hours; 31.0% had a fatty meal and 31.0% had a light meal as the last preoperative intake. The average liquid preoperative fasting time was 9.43 hours, with 46.1% of participants having water as their last intake.

When the fasting time of the children was investigated in accordance with the last meal they had eaten, it was found that the children were fed on average 10.51 hours before surgery ($n = 203$) with clear liquids, 6.27 hours ($n = 52$) with breast milk, 9.9 hours ($n = 107$) with baby food or cow's milk, 11.22 hours ($n = 100$) with toast and other, and 12.25 hours ($n = 106$) with daily food ([Table 2](#)).

65.6% of children between ages 0 and 3 years ($n = 151$) were noted by their parents as being

Table 1. The Sociodemographic Distribution of the Sample

	Number	Percentage
Age groups		
0 to 1 y	54	16.3
1 to 3 y	97	29.2
3 to 6 y	81	24.4
6 to 12 y	66	19.9
12 y and older	34	10.2
Gender		
Female	106	31.9
Male	226	68.1
Area of living		
Village	21	6.3
Town	8	2.4
District	120	36.1
City	183	55.1
Education		
Not going to school	201	60.5
Kindergarten	35	10.5
Primary school	74	22.3
High school	16	4.9
Special education	6	1.8
Sum	332	100

Download English Version:

<https://daneshyari.com/en/article/5570392>

Download Persian Version:

<https://daneshyari.com/article/5570392>

[Daneshyari.com](https://daneshyari.com)