



ELSEVIER

Contents lists available at [ScienceDirect](http://www.elsevier.com/locate/pcd)

Primary Care Diabetes

journal homepage: <http://www.elsevier.com/locate/pcd>PCDE
primary care diabetes europe

Brief report

A pilot service-evaluation examining change in HbA1c related to the prescription of internet-based education films for type 2 diabetes

S. Rice^a, H. Cranch^a, K. Littlemore^b, J. Mortimer^b, J. Platts^c,
J.W. Stephens^{d,e,*}

^a Hywel Dda University Health Board, Diabetes Centre, Prince Philip Hospital, Hywel Dda Health Board, Llanelli SA14 8QF, UK

^b eHealth Digital Media Ltd., 137, Newton Road, Swansea SA3 4ST, UK

^c Diabetes Centre, University Hospital Llandough, Cardiff and Vale University Health Board, Penarth CF63 2XX, UK

^d Diabetes Research Group, School of Medicine, Swansea University, Swansea, SA2 8PP, UK

^e Department of Diabetes & Endocrinology, ABM University Health Board, Swansea SA6 8NL, UK

ARTICLE INFO

Article history:

Received 25 November 2016

Received in revised form

15 February 2017

Accepted 16 February 2017

Available online xxx

Keywords:

Diabetes

Patient education

Internet

Films

HbA1c

ABSTRACT

We undertook a pilot service-evaluation of prescribed internet-based patient education films for patients with type 2 diabetes. The uptake was 28% and film watching was associated with a relative mean difference in HbA1c of -9.0 mmol/mol in the film watchers compared to non-watchers over a three-month period ($P = 0.0008$).

Crown Copyright © 2017 Published by Elsevier Ltd on behalf of Primary Care Diabetes Europe. All rights reserved.

1. Introduction

Providing education and knowledge to patients with type 2 diabetes is a mainstay in management. With respect to type 2 diabetes, these programmes are successful, but require infras-

tructure and funding to support their delivery. Furthermore, one criticism of published studies is the focus on participants that have attended the sessions [1,2]. The available literature recognises that the uptake of patient education courses for type 2 diabetes and other chronic conditions is low [3–6]. There is little information in the available literature reporting on alternative methods of delivering empirical education to

* Corresponding author at: Diabetes Research Group, School of Medicine, Swansea University, Swansea, Wales SA2 8PP, UK. Fax: +44 1792 703214.

E-mail address: J.W.Stephens@Swansea.ac.uk (J.W. Stephens).

<http://dx.doi.org/10.1016/j.pcd.2017.02.002>

1751-9918/Crown Copyright © 2017 Published by Elsevier Ltd on behalf of Primary Care Diabetes Europe. All rights reserved.

Table 1 – Changes in HbA1c before and after 3 months of internet-based educational films.

	Before	After	Mean difference	P-value
Non watchers (n = 49)				
HbA1c (mmol/mol)	59.1 ± 10.6	60.5 ± 13.2*	+1.7**	0.23
Watchers (n = 19)				
HbA1c (mmol/mol)	57.3 ± 16.0	49.8 ± 10.1*	-7.4**	0.03

* P = 0.0008 for HbA1c at follow-up in Watchers compared to Non-watchers.

** P = 0.01.

patients with type 2 diabetes. If this could be delivered in an economically and clinically effective manner then this might act as a stepping-stone to more structured patient education. Our aim was to undertake a three-month pilot service evaluation of prescribed internet-based patient education films for patients with type 2 diabetes within primary care.

2. Methods

2.1. Participants

This work was approved and undertaken as a service-evaluation project within Hywel Dda Health Board South Wales. An initial approach to recruitment was made to patients who had been identified with a new diagnosis of type 2 diabetes within seven primary care practices. The local practice nurses then explained the nature of the films and the process of watching these (as described below). The films were streamed directly to patients' electronic devices (home computers, smart phones, tablets) via e-mail links with one film being released each week. The study ran from the 1st of January 2016 to the 1st of October 2016. A blood sample was collected and analysed in the routine clinical chemistry laboratory at the time of the film prescription and at three months.

2.2. Video preparation and prescribing

High quality, diabetes-specific educational and motivational films funded by the Welsh Government were developed. The creation and distribution of these films was self-funded by eHealth Digital Media Ltd and subsequent films offered to the NHS on a low-cost subscription basis paid for 'per condition per patient population'.

The films focussed on diet, weight, foot care, medication and monitoring. Patients were prescribed between 8–11 films at the time of recruitment using a web based application <https://clinic.pocketmedic.org/>. Each patient consented to the evaluation by providing their name and email address to enable the application to send hyperlinks of the films by email. The application automatically creates a patient account accessed at <https://www.pocketmedic.org/by> entering their email address and a secure PIN number. There were 11 films relating to type 2 diabetes in the series each lasting 5 min. The films were made in partnership with the authored clinicians (SR, JWS). The films used in this study were as follows:- What is diabetes?; It's been a week now... Jill's story; What can I eat? Medication and Monitoring; Introducing the Eatwell Guide; Looking after your feet; Jeff's story; So What Can I Do?; Dia-

betes and Weight; Tony and Michelle both live with diabetes; Stop smoking. The content of the films allows type 2 diabetes to be explained using clear visuals. Expert patients describe their understanding of type 2 diabetes, how this affects their lives; and management with diet, exercise, medication or injectable therapies. "What can I Eat?" is addressed along with healthy eating principles. Patients share real experiences to engage others with understanding the principles of diabetes self-management in partnership with their healthcare professionals. The content of the films are based on the principles of motivational interviewing and self-determination including patient autonomy, competence and belonging [7].

2.3. Outcomes and statistical analysis

The overall watch rate was calculated as the percentage of the total films watched relative to the total number of films made available to all the patients. A paired t-test was used to compare differences in HbA1c during the follow-up period. An ANOVA was performed to examine the change in HbA1c within the group, correcting for the baseline HbA1c. A Pearson correlation was performed to examine the association between change in HbA1c and the number of films watched.

3. Results

Sixty-eight patients (30 males, 38 females) with a mean age of 68 ± 18 years were prescribed the films. Of these, 49 did not watch any (non-watchers), whilst 19 (28%) watched the films (watchers). There was considerable variation in the number of films watched (median 4; range 1–11) and the overall watch rate was 13% (82/633). There was no significant difference in age between the watchers and non-watchers (63.9 ± 11.4 vs. 65.5 ± 12.1 years, $P=0.61$). No differences were observed at baseline in HbA1c. As shown in Table 1, there was a significant mean decrease in HbA1c of 7.1 mmol/mol in the watchers. No decrease was observed in the non-watchers. Of interest, at follow-up the mean relative difference in HbA1c between the watchers and non-watchers was -9.0 mmol/mol ($P=0.0008$). An ANCOVA was consistent with this result ($P=0.0004$), demonstrating that the effect was maintained after correcting for the baseline HbA1c. As described, within the watchers there was variation in the number of films that the patients watched. Of interest there was a significant correlation between the number of films watched and the reduction in HbA1c ($r = -0.58$, $P=0.01$), (Fig. 1).

Download English Version:

<https://daneshyari.com/en/article/5571485>

Download Persian Version:

<https://daneshyari.com/article/5571485>

[Daneshyari.com](https://daneshyari.com)