



## Original article

# Handgrip strength is an independent predictor of all-cause mortality in maintenance dialysis patients



Barbara Perez Vogt, Mariana Clementoni Costa Borges, Cassiana Regina de Goés, Jacqueline Costa Teixeira Caramori\*

Faculdade de Medicina de Botucatu, UNESP Univ Estadual Paulista, Department of Clinical Medicine, Botucatu, São Paulo, Brazil

## ARTICLE INFO

## Article history:

Received 20 November 2015

Accepted 24 March 2016

## Keywords:

Dialysis

Handgrip strength

Mortality

Nutritional assessment

## SUMMARY

**Background & aims:** Muscle wasting is associated with mortality in dialysis patients. The measurement of muscle mass has some limitations, while muscle strength assessment is simple, safe and allows the recognition of patients at risk of progressing to poor outcomes related to malnutrition. The aim of this study is verify if handgrip strength (HGS) is associated with all-cause mortality in patients in maintenance haemodialysis (HD) and peritoneal dialysis (PD).

**Methods:** This was an observational retrospective cohort study which included all patients in maintenance HD and PD from July 2012 to October 2014. Patients were followed-up until June 2015.

**Results:** Two-hundred sixty five patients were enrolled (218 HD and 47 PD) and they were followed for  $13.4 \pm 7.9$  months. During the follow-up period, 53 patients (20%) have died, 36 patients (13.6%) have undergone renal transplantation, 13 patients (4.9%) have switched off dialysis method and 5 patients (1.9%) have transferred to another facility. The cut-off of HGS able to predict mortality was 22.5 kg for men and 7 kg for women. Using this cut-off to fit the Kaplan–Meier survival curve, the association of HGS with all-cause mortality for both genders was confirmed. Finally, in the multivariate analysis adjusted for demographic, clinical and nutritional variables, HGS remained significant predictor of mortality, independent of dialysis modality.

**Conclusions:** HGS cut-offs that predict mortality were 22.5 kg for men and 7 kg for women. HGS was associated with mortality independent of dialysis modality.

© 2016 Elsevier Ltd and European Society for Clinical Nutrition and Metabolism. All rights reserved.

## 1. Introduction

End-stage renal disease (ESRD) patients are at increased risk of protein energy wasting (PEW), which is characterized by muscle mass loss associated with inflammation [1]. Muscle mass wasting is highly prevalent among maintenance dialysis patients. Evidence of wasting can be noted in 18–75% of these patients, and it is an important predictor of morbidity and mortality [1]. Therefore,

markers of muscle mass and muscle function could be important predictors of outcomes in this population [2–4].

Several methods have been used for determining muscle mass in dialysis patients, i.e., dual-energy X-ray absorptiometry, bioelectrical impedance analysis, and anthropometry. However, all of these methods have some relevant disadvantages [5,6]. In this context, handgrip strength (HGS) has gained considerable attention as an indicator of nutrition status and muscle function in recent research. HGS is a validated method to assess nutritional status [7]. It is fast, safe, simple, reliable, non-invasive, painless, radiation-free, and low cost [8]. It shows high inter-rater reliability and may be useful as an early indication of malnutrition [9]. Moreover, it seems not affected by hydration status [8,10], as the other methods for muscle mass assessment.

Studies with chronic kidney disease (CKD) patients have shown HGS is associated with malnutrition, assessed by Malnutrition Inflammation Score (MIS) [11] and renal outcomes (pre-dialysis mortality or reaching ESRD) [12]. In patients on maintenance

*Abbreviations:* ESRD, end-stage renal disease; PEW, protein energy wasting; HGS, handgrip strength; CKD, chronic kidney disease; HD, haemodialysis; PD, peritoneal dialysis; CRP, C – reactive protein; BMI, body mass index; MAMC, mid-arm muscle circumference.

\* Corresponding author. Faculdade de Medicina de Botucatu, UNESP, Univ Estadual Paulista Distrito de Rubião Junior, s/n 18618970, Botucatu, São Paulo, Brazil. Tel.: +55 14 3880 1171; fax: +55 14 3882 2238.

E-mail address: [jteixeir@fmb.unesp.br](mailto:jteixeir@fmb.unesp.br) (J.C.T. Caramori).

dialysis, HGS is associated with malnutrition evaluated by Subjective Global Assessment [13,14] and mortality [4,10].

In view of the simplicity and advantages of HGS assessment in evaluating nutritional status and the significant influence of muscle wasting on mortality of patients on dialysis, the aims of this study are: to determine a HGS cut-off for men and women and verify its association with mortality in patients on maintenance dialysis.

## 2. Materials and methods

### 2.1. Study design and subjects

It was a retrospective cohort study which included prevalent haemodialysis (HD) and peritoneal dialysis (PD) patients for at least three months, treated at Hospital of Botucatu Medical School from July 2012 to August 2014. Patients younger than 18 years and those with missing nutritional assessment data were excluded. Patients were evaluated and followed until death, transplantation, switch off dialysis method, transfer to another facility or the end of follow-up in June 2015. The study protocol was approved by local research ethics committee.

Following demographic, clinical and laboratory data were obtained from medical records: gender, age, dialysis vintage, main cause of ESRD, diabetes, serum urea, creatinine, albumin and C-reactive protein (CRP).

### 2.2. Assessment of nutritional status and muscle function

Anthropometric measurements were obtained from nutritional assessment performed after HD session or during routine outpatient care for PD patients. Body weight, height, mid-arm circumference and triceps skinfold thickness were measured. From these measurements, it was calculated body mass index (BMI), mid-arm muscle circumference (MAMC) and percent standard of MAMC was obtained from the National Health and Nutrition Examination Survey percentile distribution tables [15,16].

HGS was measured during nutritional assessment, using Jamar<sup>®</sup> mechanical dynamometer with a precision of 1 kg, in the dominant hand or in the non-fistula hand if implanted. Patients were instructed to self-adjust the dynamometer so that they fit comfortably to their hand size to get the best performance and to hold the grip with maximum force in response to a voice command, with the arm extended sideways from the body with the dynamometer facing away from the body. Three measurements were performed with intervals of about 30 s between each run and the maximum value was considered.

### 2.3. Statistical analyses

Data were expressed as mean  $\pm$  standard deviation or median (first and third quartiles), and frequencies were expressed as percentage. Comparisons between genders and between survival and non-survival patients were performed using t Student's test or Mann Whitney. Frequencies were compared by qui-square test. Sensitivity and specificity analysis was performed to verify the best HGS cut-off able to predict mortality for men and women. A Receiver Operating Characteristic Curve Analysis (ROC curve) was constructed for each gender to verify the area under the curve and the significance of HGS on mortality prediction. Kaplan–Meier survival curves were fitted, and the difference between the curves was assessed by log-rank test. Cox proportional hazards analysis was used to assess independent predictors of mortality through models adjusted for variables significantly different between survival and non-survival patients. In this analysis, HG was included as a categorical variable according to the cut-off for each gender.

Statistical significance was accepted as a p-value  $<0.05$ . Statistical analysis was performed using SPSS 22.0.

## 3. Results

### 3.1. Baseline characteristics

This study enrolled 265 patients on maintenance dialysis, most of them on HD (82.3%). Patients' age ranged from 18 to 91 years, and 54% were male. Diabetic nephropathy was the main cause of ESRD (31.1%), followed by hypertensive nephrosclerosis (18.1%). Patients mean BMI was  $26.3 \pm 7.1$  kg/m<sup>2</sup> and mean HGS was  $18.7 \pm 11.2$  kg. Mean HGS for men was  $24 \pm 11.6$  kg and for women was  $12.5 \pm 6.7$  kg ( $p < 0.001$ ).

Demographic, clinical and nutritional data of entire cohort and according to gender is presented in Table 1.

Comparing patients according to dialysis methods, HD patients were majority male, with longer time of dialysis vintage and longer follow-up, and higher values of serum urea, creatinine and albumin ( $p < 0.05$ ) than PD patients. HGS and frequency of outcomes were not different between HD and PD.

### 3.2. Comparisons between survival and non-survivals patients

Age, prevalence of diabetes, dialysis vintage and CRP were higher among non-survival patients, while serum urea, creatinine and albumin were higher among survival patients (Table 2).

### 3.3. Follow-up

Patients were followed-up for a mean  $13.4 \pm 7.9$  months (minimum 0.6 and maximum 30.9 months). During this period, 53 patients (20%) have died, 36 patients (13.6%) have received renal transplantation, 13 patients (4.9%) have switched off dialysis method and 5 patients (1.9%) have been transferred to another facility.

### 3.4. ROC curve

According to ROC curves, HGS cut-off able to predict mortality were 22.5 kg for male, with 61% sensitivity and 76% specificity (AUC 0.689; CI 95% 0.575–0.803;  $p < 0.003$ ) and 7 kg for female, with 83% sensitivity and 35.7% specificity (AUC 0.615; CI 95% 0.493–0.737;  $p = 0.06$ ) (see Fig. 1).

### 3.5. Kaplan–Meier survival analysis

Survival probability analysis (Kaplan–Meier) confirmed the association of HGS with all-cause mortality for both gender (Fig. 2). The results showed statistical significance for both male (Fig. 2A) ( $p = 0.003$ ) and female (Fig. 2 B) ( $p = 0.004$ ).

### 3.6. Cox proportional hazards analysis

Models were fitted using cox proportional hazards analysis with HGS as a categorical variable, considering the cut-off for each gender (Table 3). First, crude analysis showed HGS as a significant predictor of mortality. HGS remained significant predictor of mortality even after adjustments.

## 4. Discussion

The present study showed HGS was able to identify increased risk of all-cause mortality in dialysis patients, with different cut-offs for genders. HGS is a measure widely used in clinical

Download English Version:

<https://daneshyari.com/en/article/5572034>

Download Persian Version:

<https://daneshyari.com/article/5572034>

[Daneshyari.com](https://daneshyari.com)