

Accepted Manuscript

Meaningful communication prior to death, but not presence at the time of death itself, is associated with better outcomes on measures of depression and complicated grief among bereaved family members of cancer patients

Hiroyuki Otani, M.D., Saran Yoshida, Ph.D., C.P, Tatsuya Morita, M.D., Maho Aoyama, R.N., Ph.D., Yoshiyuki Kizawa, M.D., Ph.D., Yasuo Shima, M.D., Satoru Tsuneto, M.D., Ph.D., Mitsunori Miyashita, R.N., Ph.D.

PII: S0885-3924(17)30273-7

DOI: [10.1016/j.jpainsymman.2017.07.010](https://doi.org/10.1016/j.jpainsymman.2017.07.010)

Reference: JPS 9435

To appear in: *Journal of Pain and Symptom Management*

Received Date: 16 February 2017

Revised Date: 27 May 2017

Accepted Date: 6 July 2017

Please cite this article as: Otani H, Yoshida S, Morita T, Aoyama M, Kizawa Y, Shima Y, Tsuneto S, Miyashita M, Meaningful communication prior to death, but not presence at the time of death itself, is associated with better outcomes on measures of depression and complicated grief among bereaved family members of cancer patients, *Journal of Pain and Symptom Management* (2017), doi: 10.1016/j.jpainsymman.2017.07.010.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Meaningful communication prior to death, but not presence at the time of death itself, is associated with better outcomes on measures of depression and complicated grief among bereaved family members of cancer patients

Running title: Grief and family presence at patient's death

Hiroyuki Otani, M.D.

Department of Palliative Care Team, and Palliative and Supportive Care, National Kyushu Cancer Center

3-1-1 Notame, Mitami-ku, Fukuoka, 811-1395, Japan.

Saran Yoshida, Ph.D., C.P.

Department of Clinical Psychology, Tohoku University Graduate School of Education

27-1, Kawauchi, Aoba-ku, Sendai, Miyagi 980-8576, Japan

Tatsuya Morita, M.D.

Department of Palliative and Supportive Care, Palliative Care Team, Seirei Mikatahara General Hospital

3453 Mikatahara-cho, Kita-ku, Hamamatsu, Shizuoka 433-8558, Japan

Maho Aoyama, R.N., Ph.D.

Department of Community Health, Health Sciences, Tohoku University Graduate School of Medicine

2-1 Seiryomachi, Aoba-ku, Sendai, Miyagi 980-8575, Japan

Yoshiyuki Kizawa, M.D., Ph.D.

Department of Palliative Medicine, Kobe University Graduate School of Medicine

7-5-1, Kusunoki-cho, Chuo-ku, Kobe, Hyogo 650-0017, Japan

Download English Version:

<https://daneshyari.com/en/article/5578191>

Download Persian Version:

<https://daneshyari.com/article/5578191>

[Daneshyari.com](https://daneshyari.com)