

Accepted Manuscript

The conversion ratio from intravenous hydromorphone to oral opioids in cancer patients

Akhila Reddy, MD, Marieberta Vidal, MD, Saneese Stephen, PA, Karen Baumgartner, APRN, Sara Dost, MD, Ann Nguyen, MD, Yvonne Heung, MD, Simeon Kwan, DO, Angelique Wong, MD, Imelda Pangemanan, APRN, Ahsan Azhar, MD, Supakarn Tayjasanant, MD, Edenmae Rodriguez, PharmD, Jessica Waletich, PharmD, Kyu-Hyoung Lim, MD, Jimin Wu, MS, Diane Liu, MS, Janet Williams, MPH, Sriram Yennurajalingam, MD, Eduardo Bruera, MD



PII: S0885-3924(17)30264-6

DOI: [10.1016/j.jpainsymman.2017.07.001](https://doi.org/10.1016/j.jpainsymman.2017.07.001)

Reference: JPS 9426

To appear in: *Journal of Pain and Symptom Management*

Received Date: 14 March 2017

Revised Date: 14 June 2017

Accepted Date: 7 July 2017

Please cite this article as: Reddy A, Vidal M, Stephen S, Baumgartner K, Dost S, Nguyen A, Heung Y, Kwan S, Wong A, Pangemanan I, Azhar A, Tayjasanant S, Rodriguez E, Waletich J, Lim K-H, Wu J, Liu D, Williams J, Yennurajalingam S, Bruera E, The conversion ratio from intravenous hydromorphone to oral opioids in cancer patients, *Journal of Pain and Symptom Management* (2017), doi: 10.1016/j.jpainsymman.2017.07.001.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The conversion ratio from intravenous hydromorphone to oral opioids in cancer patients

Akhila Reddy, MD^{1*}; Marieberta Vidal, MD^{1*}; Saneese Stephen, PA¹; Karen Baumgartner, APRN¹; Sara Dost, MD¹; Ann Nguyen, MD¹; Yvonne Heung, MD¹; Simeon Kwan, DO¹; Angelique Wong, MD¹; Imelda Pangemanan, APRN¹; Ahsan Azhar, MD¹; Supakarn Tayjasanant, MD²; Edenmae Rodriguez, PharmD¹; Jessica Waletich, PharmD¹; Kyu-Hyoung Lim, MD³; Jimin Wu, MS⁴; Diane Liu, MS⁴; Janet Williams, MPH¹; Sriram Yennurajalingam, MD¹; and Eduardo Bruera, MD¹

¹*Department of Palliative Care and Rehabilitation Medicine and* ⁴*Department of Biostatistics The University of Texas MD Anderson Cancer Center, Houston, TX, USA*

²*Siriraj Palliative Care Center, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok, Thailand*

³*Department of Internal Medicine, Kangwon National University School of Medicine, South Korea*

*AR and MV contributed equally to the study.

Corresponding author

Akhila Reddy, MD

Department of Palliative, Rehabilitation, and Integrative Medicine; Unit 1414

The University of Texas MD Anderson Cancer

1515 Holcombe Blvd.

Houston, TX USA, 77030

Phone: 713-792-7567; Fax: 713-792-6092

E-mail: asreddy@mdanderson.org

Key words: Cancer patients, opioid rotation, Intravenous hydromorphone, morphine equivalent daily dose, conversion ratio, and cancer pain

Running title: Conversion from IV hydromorphone to oral opioids

Download English Version:

<https://daneshyari.com/en/article/5578192>

Download Persian Version:

<https://daneshyari.com/article/5578192>

[Daneshyari.com](https://daneshyari.com)