



Continuous deep sedation: A proposal for performing more rigorous empirical research.

Tatsuya Morita, M.D.

Department of Palliative and Supportive Care, Palliative Care Team, Seirei Mikatahara General Hospital
3453 Mikatahara-cho, Kita-ku, Hamamatsu, Shizuoka 433-8558, Japan

Kengo Imai, M.D.

Seirei Hospice, Seirei Mikatahara General Hospital
3453 Mikatahara-cho, Kita-ku, Hamamatsu, Shizuoka 433-8558, Japan

Naosuke Yokomichi, M.D., Ph.D.

Seirei Hospice, Seirei Mikatahara General Hospital
3453 Mikatahara-cho, Kita-ku, Hamamatsu, Shizuoka 433-8558, Japan

Masanori Mori, M.D.

Palliative Care Team, Seirei Mikatahara General Hospital
3453 Mikatahara-cho, Kita-ku, Hamamatsu, Shizuoka 433-8558, Japan

Yoshiyuki Kizawa M.D., Ph.D.

Department of Palliative Medicine, Kobe University Graduate School of Medicine
7-5-1, Kusunoki-cho, Chuo-ku, Kobe, Hyogo 650-0017, Japan

Satoru Tsuneto, M.D., Ph.D.

Palliative Care Center, Department of Palliative Medicine, Kyoto University Hospital
54 Kawaharacho, Shogoin, Sakyo-ku, Kyoto 606-8507, Japan

Words 2,167, 3 Tables, 2 Figures

Download English Version:

<https://daneshyari.com/en/article/5578533>

Download Persian Version:

<https://daneshyari.com/article/5578533>

[Daneshyari.com](https://daneshyari.com)