

# Accepted Manuscript

Low Pretreatment Neutrophil-To-Lymphocyte Ratio Predicts Good Outcome in Patients treated with Neoadjuvant Chemotherapy Prior to Radical Cystectomy for Muscle Invasive Bladder Cancer

Oscar Buisan, Anna Orsola, Joan Areal, Albert Font, Mario Oliveira, Roberto Martinez, Luis Ibarz

PII: S1558-7673(16)30130-6

DOI: [10.1016/j.clgc.2016.05.004](https://doi.org/10.1016/j.clgc.2016.05.004)


Reference: CLGC 613

To appear in: *Clinical Genitourinary Cancer*

Received Date: 23 February 2016

Revised Date: 23 April 2016

Accepted Date: 18 May 2016



Please cite this article as: Buisan O, Orsola A, Areal J, Font A, Oliveira M, Martinez R, Ibarz L, Low Pretreatment Neutrophil-To-Lymphocyte Ratio Predicts Good Outcome in Patients treated with Neoadjuvant Chemotherapy Prior to Radical Cystectomy for Muscle Invasive Bladder Cancer, *Clinical Genitourinary Cancer* (2016), doi: 10.1016/j.clgc.2016.05.004.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Low Pretreatment Neutrophil-To-Lymphocyte Ratio Predicts Good Outcome in Patients treated with Neoadjuvant Chemotherapy Prior to Radical Cystectomy for Muscle Invasive Bladder Cancer**

Oscar Buisan<sup>1</sup>, Anna Orsola<sup>2</sup>, Joan Areal<sup>1</sup>, Albert Font<sup>3</sup>,  
Mario Oliveira<sup>1</sup>, Roberto Martinez<sup>1</sup>, Luis Ibarz<sup>1</sup>

<sup>1</sup>Urology Department *Hospital Universitari Germans Trias i Pujol* and Autonomous University of Barcelona, Barcelona, Spain,

<sup>2</sup>Bladder Cancer Center, Dana-Farber Cancer Institute, Boston, Massachusetts, USA,

<sup>3</sup>Medical Oncology Service, Department of Medicine, Catalan Institute of Oncology, *Hospital Universitari Germans Trias i Pujol* and Autonomous University of Barcelona, Barcelona, Spain

**Runninghead:** neutrophil-to-lymphocyte ratio in neoadjuvant chemotherapy treated bladder cancer.

**Keywords:** Bladder cancer; Neoadjuvant Therapy; Neutrophil lymphocyte ratio; Radical cystectomy; Urothelial cancer

**Word count:** abstract = 256, text = 2469

**Number of:** tables = 4, figures = 1 (+ 2 supplementary figure)

**Relevant disclosures and conflict of interest:** All coauthors: none

**Abbreviations**

MIBC – muscle-invasive-bladder cancer

NAC – neoadjuvant chemotherapy

NLR – neutrophil-lymphocyte ratio

RC – radical cystectomy

PR – pathologic response

TURBT - transurethral resection of bladder tumor

Download English Version:

<https://daneshyari.com/en/article/5581144>

Download Persian Version:

<https://daneshyari.com/article/5581144>

[Daneshyari.com](https://daneshyari.com)