



## Four Dimensional Transesophageal Echocardiography guided AngioVac Debulking of a Tricuspid Valve Vegetation

Adam A. Dalia MD,MBA<sup>1</sup>, Daniel Bamira,MD<sup>2</sup>, Mazen Albaghdadi,MD<sup>3</sup>, Michael Essandoh,MD<sup>4</sup>, Kenneth Rosenfield,MD<sup>3</sup>, David Dudzinski,MD<sup>2</sup>

1. Department of Anesthesiology, Pain Medicine, and Critical Care, The Massachusetts General Hospital, Harvard Medical School, Boston, MA
2. Department of Cardiology, The Massachusetts General Hospital, Harvard Medical School, Boston, MA
3. Department of Interventional Cardiology, The Massachusetts General Hospital, Harvard Medical School, Boston, MA
4. Department of Anesthesiology, The Ohio State University Wexner Medical Center, Columbus, OH

Corresponding Author:

Adam A. Dalia, MD, MBA

CardioThoracic Anesthesiology Fellow

Department of Anesthesia, Critical Care and Pain Medicine

The Massachusetts General Hospital, Harvard Medical School

55 Fruit St. Boston, MA 02114

aadalia@mgh.harvard.edu

Cell: (513) 375-9015

Keywords:

TEE; echocardiography; 4D TEE; Live 3D TEE; AngioVac; Vortex; Tricuspid Valve Vegetation; TEE Guided Catheter Based procedures; Intraprocedural TEE guidance

### Introduction:

There is scarce literature regarding the use of large volume aspiration/suction catheters (Vortex AngioVac suction device; AngioDynamics, Latham, NY) for tricuspid valve vegetation

Download English Version:

<https://daneshyari.com/en/article/5582275>

Download Persian Version:

<https://daneshyari.com/article/5582275>

[Daneshyari.com](https://daneshyari.com)