


# Author's Accepted Manuscript

## Vector Flow Mapping and Impaired Left Ventricular Flow by Alfieri Stitch

Koichi Akiyama, Keiichi Itatani, Yoshifumi Naito, Mao Kinoshita, Masaru Shimizu, Saeko Hamaoka, Hiroaki Yasumoto, Hideya Kato, Yasufumi Nakajima, Satoshi Numata, Hitoshi Yaku, Teiji Sawa



PII: S1053-0770(16)30277-4  
DOI: <http://dx.doi.org/10.1053/j.jvca.2016.07.013>  
Reference: YJCAN3759

To appear in: *Journal of Cardiothoracic and Vascular Anesthesia*

Received date: 10 January 2016

Cite this article as: Koichi Akiyama, Keiichi Itatani, Yoshifumi Naito, Mao Kinoshita, Masaru Shimizu, Saeko Hamaoka, Hiroaki Yasumoto, Hideya Kato, Yasufumi Nakajima, Satoshi Numata, Hitoshi Yaku and Teiji Sawa, Vector Flow Mapping and Impaired Left Ventricular Flow by Alfieri Stitch, *Journal of Cardiothoracic and Vascular Anesthesia*, <http://dx.doi.org/10.1053/j.jvca.2016.07.013>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Vector Flow Mapping and Impaired Left Ventricular Flow by Alfieri Stitch**

Koichi Akiyama, MD; Keiichi Itatani, MD, PhD; Yoshifumi Naito, MD; Mao Kinoshita MD;  
Masaru Shimizu MD; Saeko Hamaoka MD; Hiroaki Yasumoto MD; Hideya Kato MD;  
Yasufumi Nakajima, MD, PhD; Satoshi Numata, MD, PhD; Hitoshi Yaku, MD, PhD; Teiji  
Sawa, MD, PhD

From the Departments of Anesthesiology (K.A., Y.N., M.K., M.S., S.H., H.Y., H.K., Y.N.,  
T.S.) and Cardiovascular Surgery (K.I., S.N., H.Y.), Kyoto Prefectural University of  
Medicine, Kyoto, Japan, and Department of Anesthesiology, Kansai Medical University,  
Hirakata, Japan (Y.N.)

Vector Flow Mapping and Alfieri Stitch

**Correspondence to Koichi Akiyama MD.**

Department of Anesthesiology, Kyoto Prefectural University of Medicine.  
465, Kajii Cho, Hirokoji Agaru, Kawaramachi Street,  
Kamigyo Ward, Kyoto City, Kyoto Prefecture 602-8566, Japan.  
Telephone: 075-251-5633; Fax: 075-251-5843; E-mail: [kanaslik@koto.kpu-m.ac.jp](mailto:kanaslik@koto.kpu-m.ac.jp)

**Acknowledgments**

The authors would like to thank Enago ([www.enago.jp](http://www.enago.jp)) for the English language review.

**Ethics approval**

Informed consent was obtained from the patient for the present clinical study.

**Funding**

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

**Conflicts of interest:** None

Download English Version:

<https://daneshyari.com/en/article/5582360>

Download Persian Version:

<https://daneshyari.com/article/5582360>

[Daneshyari.com](https://daneshyari.com)