Error without trials: Safe SpO2 threshold levels may not be derivable from SpO2-PaO2 relationships

Gareth L. Jones, Samiran Ray, Padmanabhan Ramnarayan, Mark J. Peters

PII: S0883-9441(17)30735-9
DOI: doi: 10.1016/j.jcrc.2017.05.013
Reference: YJCRC 52525

To appear in:

Revised date: ###REVISEDDATE###
Accepted date: ###ACCEPTEDDATE###

Please cite this article as: Gareth L. Jones, Samiran Ray, Padmanabhan Ramnarayan, Mark J. Peters , Error without trials: Safe SpO2 threshold levels may not be derivable from SpO2-PaO2 relationships, (2017), doi: 10.1016/j.jcrc.2017.05.013

Error without trials: Safe SpO_2 threshold levels may not be derivable from SpO_2 - PaO_2 relationships.

Gareth L Jones [1, 2.] Samiran Ray [1, 2..] Padmanabhan Ramnarayan[3.] Mark J Peters [1, 2]

Author Affiliations:

1. Respiratory Critical Care and Anaesthesia Unit, UCL Great Ormond Street Institute of Child Health, London, United Kingdom.
2. Paediatric Intensive Care Unit, Great Ormond Street Hospital, London, United Kingdom.
3. Children's Acute Transport service, Great Ormond Street Hospital, London, United Kingdom.

Corresponding author:

Gareth L Jones

Address: Paediatric Intensive Care Unit, Great Ormond Street Hospital, London, United Kingdom.

Email address: g.jones@ucl.ac.uk

Funding:

This work was undertaken at Great Ormond Street Hospital/UCL Institute of Child Health, which received a proportion of funding from the Department of Health's NIHR Biomedical Research Centre's funding scheme. Gareth Jones is a NIHR funded Academic Clinical Fellow.

Key words:

Hyperoxia, pulse oximetry, saturation, critical care, child