

Adjusting FRAX® on TBS for identification of subjects at high risk of fractures

G. Couraud, C. Souffir, E. Gaigneux, S. Kolta, C. Roux, K. Briot

PII: S8756-3282(17)30166-7
DOI: doi: 10.1016/j.bone.2017.05.005
Reference: BON 11314

To appear in: *Bone*

Received date: 17 January 2017
Revised date: 17 April 2017
Accepted date: 3 May 2017

Please cite this article as: G. Couraud, C. Souffir, E. Gaigneux, S. Kolta, C. Roux, K. Briot , Adjusting FRAX® on TBS for identification of subjects at high risk of fractures, *Bone* (2017), doi: 10.1016/j.bone.2017.05.005

Adjusting FRAX® on TBS for identification of subjects at high risk of fractures

G. Couraud[1], C. Souffir[1, 2], E. Gaigneux[1], S. Kolta[1, 2], C. Roux[1, 2,3], K. Briot[1,2]

1. Department of Rheumatology, Cochin Hospital, Assistance Publique- Hôpitaux de Paris, Paris, France.
2. INSERM U1153, Paris, France.
3. Paris-Descartes University, Paris, France.

For correspondence:

Karine Briot, MD, PhD

Hôpital Cochin, Service de rhumatologie

27 rue du Faubourg St Jacques, 75014 Paris, France .

Email: karine.briot@aphp.fr

Phone number: +33 1 58 41 25 84

Fax number: +33 1 58 41 13 70

Key words: fracture, FRAX, TBS, fracture liaison service