

Accepted Manuscript

Neutrophil-to-lymphocyte ratio is associated with diabetic peripheral neuropathy in type 2 diabetes patients

Siyong Liu, Hangping Zheng, Xiaoming Zhu, Fei Mao, Shuo Zhang, Hongli Shi, Yiming Li, Bin Lu

PII: S0168-8227(17)30074-8

DOI: <http://dx.doi.org/10.1016/j.diabres.2017.05.008>

Reference: DIAB 6965


To appear in: *Diabetes Research and Clinical Practice*

Received Date: 13 January 2017

Accepted Date: 8 May 2017

Please cite this article as: S. Liu, H. Zheng, X. Zhu, F. Mao, S. Zhang, H. Shi, Y. Li, B. Lu, Neutrophil-to-lymphocyte ratio is associated with diabetic peripheral neuropathy in type 2 diabetes patients, *Diabetes Research and Clinical Practice* (2017), doi: <http://dx.doi.org/10.1016/j.diabres.2017.05.008>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Neutrophil-to-lymphocyte ratio is associated with
diabetic peripheral neuropathy in type 2 diabetes
patients**

Siying Liu*, Hangping Zheng*, Xiaoming Zhu, Fei Mao, Shuo Zhang, Hongli Shi,

Yiming Li, Bin Lu

Department of Endocrinology and Metabolism, Huashan Hospital, Fudan

University

12# Middle Wulumuqi Road, Shanghai, China, 200040

* These authors contributed equally to this work

Corresponding author: Yiming Li: yimingli@fudan.edu.cn

And Bin Lu: doctor_lubin@126.com

Download English Version:

<https://daneshyari.com/en/article/5587048>

Download Persian Version:

<https://daneshyari.com/article/5587048>

[Daneshyari.com](https://daneshyari.com)