



Contents lists available at ScienceDirect

## Journal of Diabetes and Its Complications

journal homepage: [WWW.JDCJOURNAL.COM](http://WWW.JDCJOURNAL.COM)

## Leptin is associated with cardiopulmonary fitness independent of body-mass index and insulin sensitivity in adolescents with type 1 diabetes: a brief report from the EMERALD study

Petter Bjornstad<sup>a,b,\*</sup>, Melanie Cree-Green<sup>a,c</sup>, Amy Baumgartner<sup>a</sup>, Gregory Coe<sup>a</sup>, Yesenia Garcia Reyes<sup>a</sup>, Michal Schafer<sup>d</sup>, Laura Pyle<sup>a</sup>, Judith G. Regensteiner<sup>c,e</sup>, Jane E.B. Reusch<sup>c,f</sup>, Kristen J. Nadeau<sup>a,c</sup>

<sup>a</sup> Department of Pediatric Endocrinology, University of Colorado School of Medicine, Aurora, CO, United States

<sup>b</sup> Barbara Davis Center for Childhood Diabetes, University of Colorado Denver, Aurora, CO

<sup>c</sup> Center for Women's Health Research, Divisions of General Internal Medicine and Cardiology, University of Colorado School of Medicine, Aurora, CO, United States

<sup>d</sup> Division of Pediatric Cardiology, University of Colorado School of Medicine, Aurora, CO, United States

<sup>e</sup> Divisions of General Internal Medicine and Cardiology, University of Colorado School of Medicine, Aurora, CO, United States

<sup>f</sup> Division of Endocrinology, Veterans Administration Hospital, University of Colorado School of Medicine, Aurora, CO, United States

## ARTICLE INFO

## Article history:

Received 21 December 2016

Received in revised form 28 January 2017

Accepted 13 February 2017

Available online xxx

## Keywords:

Leptin

Type 1 diabetes

Cardiopulmonary fitness

Adiposity

Insulin sensitivity

## ABSTRACT

Cardiopulmonary fitness is decreased in type 1 diabetes for reasons that are incompletely understood. In this study, leptin was associated with exercise capacity independent of insulin sensitivity (IS) and body mass index (BMI), suggesting that leptin may relate to cardiopulmonary fitness by mechanisms beyond IS and/or obesity.

© 2016 Published by Elsevier Inc.

## 1. Introduction

Cardiovascular disease (CVD) is the leading cause of morbidity and mortality in type 1 diabetes (T1D)<sup>1,2</sup> with abnormalities measurable in adolescence.<sup>3</sup> In T1D youth we have reported an inverse relationship between insulin sensitivity (IS) and peak exercise capacity (VO<sub>2</sub>peak).<sup>4</sup> Adipocyte dysfunction is linked to insulin resistance,<sup>5</sup> which is an important factor in CVD and cardiorenal disease in T1D.<sup>6</sup> Leptin, an adipocytokine, is increased in obesity, and induces metabolic and cardiovascular actions that may contribute to CVD. Leptin is also associated with IS<sup>7</sup> and we previously demonstrated that higher IS relates to better peak exercise capacity (VO<sub>2</sub>peak).<sup>4</sup> Adiposopathy, as assessed by low plasma adiponectin/leptin ratio has also been proposed to contribute to cardiovascular disease.<sup>8</sup> However, it is unknown whether leptin or adiponectin/leptin ratio are related to cardiopulmonary fitness itself, and whether this relationship is dependent or independent of adiposity and/or IS in youth with T1D. Accordingly, we tested the hypothesis that circulating leptin concentration is associated with VO<sub>2</sub>peak independent of IS and BMI, in youth with T1D.

None of the authors have any conflicts of interest to disclose.

\* Corresponding author at: 13123 East 16th Ave Box B265, Aurora, CO 80045. Tel.: +1 720 777 4659; fax: +1 720 777 7301.

E-mail address: [Petter.Bjornstad@childrenscolorado.org](mailto:Petter.Bjornstad@childrenscolorado.org) (P. Bjornstad).

## 2. Methods

## 2.1. Participants

Forty-one pubertal adolescents ages 12 to 21 years were included from the **E**ffects of **M**etformin on Cardiovascula**R** Function in **A**do**L**escents with **T**ype 1 **D**iabetes (EMERALD) study. Participants were recruited from the Barbara Davis Center and private practices with advertisements. The study was approved by the University of Colorado Denver Institutional Review Board, with appropriate consent and assent obtained. Pubertal development was assessed by a single pediatric endocrinologist (KJN) as previously described.<sup>4,9</sup> After 5 min of supine rest, 3 blood pressure measurements were obtained and averaged using a DynaPulse 5200A (Pulse Metric, San Diego, California). Height was measured to the nearest 0.1 cm using a wall-mounted stadiometer; weight was measured to the nearest 0.1 kg using a Detecto scale (Detecto, Webb City, Missouri).

Laboratory testing was performed after an inpatient 12-h fast with an overnight intravenous insulin infusion to normalize glycemia.<sup>4,9</sup> Leptin and adiponectin were measured via radioimmunoassay (Millipore, Billerica MA). Other fasting laboratory evaluations included: total cholesterol, low-density lipoprotein cholesterol (LDL-C), high-density lipoprotein (HDL-C) cholesterol, triglycerides, glucose and HbA1c (DCCT-calibrated);

<http://dx.doi.org/10.1016/j.jdiacomp.2017.02.019>  
1056-8727/© 2016 Published by Elsevier Inc.

Please cite this article as: Bjornstad P, et al. Leptin is associated with cardiopulmonary fitness independent of body-mass index and insulin sensitivity in adolescents with type .... *Journal of Diabetes and Its Complications* (2017), <http://dx.doi.org/10.1016/j.jdiacomp.2017.02.019>

**Table 1**  
Clinical characteristics of adolescents with type 1 diabetes across tertiles of leptin.

Variables	Low leptin tertile (<10.6 ng/mL) [n = 13]	Mid leptin tertile (10.6–26 ng/mL) [n = 13]	High leptin tertile (>26 ng/mL) [n = 14]	p-value
Age (years)	16.9 ± 1.6	17.4 ± 2.4	15.5 ± 2.1*	0.051
Weight (kg)	81.8 ± 41.4	72.5 ± 13.7	79.4 ± 13.7	0.64
Height (cm)	158.7 ± 36.3	165.3 ± 7.2	162. ± 16.8	0.77
Tanner stage	5 (4–5)	5 (5–5)	5 (5–5)	0.53
Female (%)	15%	62%	79%	0.003
Fasting glucose (mg/dL)	114 ± 16	110 ± 19	112 ± 17	0.80
HbA1c (%)	9.2 ± 1.6	8.3 ± 0.9	8.8 ± 1.7	0.30
BMI (%tile)	59 ± 26	81 ± 21‡	92 ± 8‡	0.0005
Resting HR (min <sup>-1</sup> )	65 ± 11	72 ± 14	76 ± 8	0.01
Rate pressure product (mm Hg * min <sup>-1</sup> )	7614 ± 1461	8887 ± 1352	9407 ± 1492	0.008
SBP (mm Hg)	117 ± 8	122 ± 10	124 ± 10	0.19
DBP (mm Hg)	68 ± 6	74 ± 5	74 ± 8	0.03
SBP (%tile)	51 ± 26	77 ± 19‡	80 ± 20‡	0.002
DBP (%tile)	48 ± 20	74 ± 14‡	72 ± 20‡	0.001
LDL-C (mg/dL)	78 ± 18	81 ± 20	94 ± 17‡	0.07
HDL-C (mg/dL)	43 ± 9	46 ± 10	48 ± 10	0.34
Total cholesterol (mg/dL)	136 ± 24	142 ± 24	160 ± 27	0.04
Triglycerides (mg/dL)	74 ± 26	73 ± 27	90 ± 52	0.42
Lean mass (kg)	49.8 ± 9.0	45.7 ± 8.5	45.9 ± 7.6†	<0.0001
Fat mass (kg)	15.5 ± 5.4	24.9 ± 5.9	31.3 ± 6.9	0.37
Leptin (ng/mL)	4.6 ± 2.7	17.1 ± 4.9	39.2 ± 12.0‡	<0.0001
Adiponectin	9.4 ± 4.4	11.8 ± 4.5	10.6 ± 5.6	0.47
Adiponectin/Leptin ratio <sup>‡</sup>	2.3 (1.6–3.2)	0.7 (0.5–1.0)	0.2 (0.2–0.3) †	<0.0001
IS (mg/kg/min)	9.5 ± 3.1	7.2 ± 3.3	4.5 ± 2.1†	0.0007
IS (mg/lean kg/min)	12.8 ± 4.1	12.3 ± 4.6	7.8 ± 3.6†	0.009
Average insulin at steady state	180.2 ± 48.3	156.5 ± 26.5	223.0 ± 89.8*	0.03
IS (mg/kg/mn)/steady state insulin	0.05 ± 0.02	0.05 ± 0.03	0.02 ± 0.01	0.0006
IS (mg/lean kg/mn)/steady state insulin	0.07 ± 0.02	0.09 ± 0.04	0.04 ± 0.03	0.001
VO <sub>2</sub> peak (mL/kg/min)	33.5 ± 6.7	22.9 ± 3.4‡	20.9 ± 3.9‡	<0.0001
VO <sub>2</sub> peak (mL/lean kg/min)	44.6 ± 6.5	36.3 ± 6.9‡	35.9 ± 6.5‡	0.002

\* p < 0.05 compared to mid tertile.

† p < 0.05 compared to mid and low tertiles.

‡ p < 0.05 compared to low tertile.

‡ Geometric means and 95% CI.

assays were performed by standard methods in the CTRC laboratory. Body composition was assessed dual-energy x-ray absorptiometry (Hologic, Waltham, MA).<sup>10</sup> IS (glucose infusion rate [GIR] in mg/kg/min and mg/lean kg/min) was calculated from a hyperinsulinemic euglycemic clamp (80 mU/m<sup>2</sup>/min insulin) the morning following the inpatient stay.<sup>4,9</sup>

VO<sub>2</sub>peak was assessed using a graded bicycle protocol to exhaustion as previously reported.<sup>4</sup> Oxygen consumption (VO<sub>2</sub>), carbon dioxide production (VCO<sub>2</sub>) and minute ventilation (VE) were measured breath-by-breath, at rest and during exercise. Respiratory exchange ratio (RER) was calculated as VCO<sub>2</sub>/VO<sub>2</sub>. To quantify the internal cardiac workload response, we calculated the rate pressure product [(RPP) = heart rate (HR) \* systolic blood pressure (SBP)].

Analyses were performed in SAS (version 9.4; SAS Institute, Cary, NC). Variables were checked for the distributional assumption of normality. ANOVA was used to test the mean of the dependent variable across tertiles of leptin. Univariable and multivariable linear regression models were employed to examine the relationships between leptin, VO<sub>2</sub>peak, unadjusted and adjusted for age, sex, BMI and IS. Linear regression results are presented as β estimate ± standard errors (SE). Significance is an α-level < 0.05.

### 3. Results

In youth with T1D we evaluated the relationships between leptin concentration, adiponectin and adiponectin/leptin ratio and VO<sub>2</sub>peak. Stratifying data by tertiles of leptin, participants in the high leptin group had significantly lower IS and VO<sub>2</sub>peak and higher blood pressure percentiles (Table 1). Leptin concentrations were significantly higher in females and individuals with elevated BMI (data not shown). After adjusting for sex and BMI or sex and IS, the relationship between leptin and VO<sub>2</sub>peak remained (Fig. 1). HbA1c and LDL-C did

not differ by leptin tertiles. Leptin correlated negatively with VO<sub>2</sub>peak and positively with RPP which remained significant after adjusting for age, sex, BMI and IS (Table 2). Whereas adiponectin concentration was not associated with VO<sub>2</sub>peak or RPP in univariable or multivariable models, adiponectin/leptin ratio was positively associated with cardiopulmonary fitness and inversely associated with RPR (Table 2).

### 4. Discussion

We found that in T1D, elevated leptin may relate to decreased cardiopulmonary fitness by mechanisms other than IS and/or obesity alone. CVD-related morbidity and mortality are markedly increased in individuals with T1D,<sup>1,2</sup> with 17 years of life lost when T1D is diagnosed at age 10.<sup>11</sup> Obesity is an increasingly prevalent problem in youth with T1D<sup>12</sup> and linked to CVD development.<sup>13</sup> Obesity induces a progressive resistance to the anorexigenic effects of leptin, which promotes further weight gain and metabolic dysfunction.<sup>14</sup> However, not all tissues are prone to leptin resistance.<sup>15</sup> The pathophysiology underlying the relationship between elevated leptin and poor cardiopulmonary fitness remains unclear. Huby et al. demonstrated that leptin contributes to CVD by upregulating aldosterone synthesis leading to increased blood pressures, and by promoting endothelial dysfunction and the expression of profibrotic markers in the heart.<sup>15</sup> Other possible mechanisms include adiposopathy<sup>16</sup> and leptin related mitochondrial dysfunction,<sup>17</sup> but these pathways have to our knowledge not been examined in T1D.

The literature is conflicting regarding whether circulating leptin concentrations are affected by T1D status,<sup>18–22</sup> and measurements of leptin in relation to VO<sub>2</sub>peak in T1D youths have to our knowledge never previously been reported.

We previously demonstrated reduced VO<sub>2</sub>peak, cardiac and vascular dysfunction in otherwise healthy, non-obese adolescents

Download English Version:

<https://daneshyari.com/en/article/5588194>

Download Persian Version:

<https://daneshyari.com/article/5588194>

[Daneshyari.com](https://daneshyari.com)