

## Accepted Manuscript

Glucose-mediated inactivation of AMP-activated protein kinase reduces the levels of L-type amino acid transporter 1 mRNA in C2C12 cells

Yu Yamamoto, Ran Sawa, Ikumi Wake, Ayaka Morimoto, Yasuhiko Okimura

PII: S0271-5317(17)30457-8  
DOI: doi: [10.1016/j.nutres.2017.08.003](https://doi.org/10.1016/j.nutres.2017.08.003)  
Reference: NTR 7791

To appear in: *Nutrition Research*

Received date: 20 May 2017  
Revised date: 5 August 2017  
Accepted date: 22 August 2017



Please cite this article as: Yamamoto Yu, Sawa Ran, Wake Ikumi, Morimoto Ayaka, Okimura Yasuhiko, Glucose-mediated inactivation of AMP-activated protein kinase reduces the levels of L-type amino acid transporter 1 mRNA in C2C12 cells, *Nutrition Research* (2017), doi: [10.1016/j.nutres.2017.08.003](https://doi.org/10.1016/j.nutres.2017.08.003)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Glucose-mediated inactivation of AMP-activated protein kinase reduces the levels of  
L-type amino acid transporter 1 mRNA in C2C12 cells**

Yu Yamamoto, Ran Sawa, Ikumi Wake, Ayaka Morimoto, Yasuhiko Okimura\*

Department of Nutrition and Food Science, Kobe Women's University Graduate School of  
Life Sciences, 2-1 Higashisuma-aoyama, Suma-ku, Kobe, 654-8585, Japan

\* Corresponding author: Phone: +81-78-737-2429; fax: +81-78-737-2429.

E-mail: y-okimura@suma.kobe-wu.ac.jp (YO)

Download English Version:

<https://daneshyari.com/en/article/5588577>

Download Persian Version:

<https://daneshyari.com/article/5588577>

[Daneshyari.com](https://daneshyari.com)