

Accepted Manuscript

De novo comparative transcriptome analysis of genes involved in fruit morphology of pumpkin cultivars with extreme size difference and development of EST-SSR markers

Aliki Xanthopoulou, Ioannis Ganopoulos, Fotis Psomopoulos, Maria Manioudaki, Theodoros Moysiadis, Aliki Kapazoglou, Maslin Osathanunkul, Sofia Michailidou, Apostolos Kalivas, Athanasios Tsaftaris, Irini Nianiou-Obeidat, Panagiotis Madesis



PII: S0378-1119(17)30286-X
DOI: doi: [10.1016/j.gene.2017.04.035](https://doi.org/10.1016/j.gene.2017.04.035)
Reference: GENE 41884
To appear in: *Gene*
Received date: 3 February 2017
Revised date: 15 April 2017
Accepted date: 19 April 2017

Please cite this article as: Aliki Xanthopoulou, Ioannis Ganopoulos, Fotis Psomopoulos, Maria Manioudaki, Theodoros Moysiadis, Aliki Kapazoglou, Maslin Osathanunkul, Sofia Michailidou, Apostolos Kalivas, Athanasios Tsaftaris, Irini Nianiou-Obeidat, Panagiotis Madesis , De novo comparative transcriptome analysis of genes involved in fruit morphology of pumpkin cultivars with extreme size difference and development of EST-SSR markers. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. *Gene*(2017), doi: [10.1016/j.gene.2017.04.035](https://doi.org/10.1016/j.gene.2017.04.035)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

***De novo* comparative transcriptome analysis of genes involved in fruit morphology of pumpkin cultivars with extreme size difference and development of EST-SSR markers**

Aliki Xanthopoulou^{1,2}, Ioannis Ganopoulos⁶, Fotis Psomopoulos³, Maria Manioudaki⁴, Theodoros Moysiadis¹, Aliki Kapazoglou¹, Maslin Osathanunkul⁵, Sofia Michailidou¹, Apostolos Kalivas⁶, Athanasios Tsaftaris^{1,2}, Irini Nianiou-Obeidat^{2#}, Panagiotis Madesis^{1#}

¹Institute of Applied Biosciences, CERTH, Thessaloniki, 570 01, Greece,

²Lab of Genetics and Plant Breeding, School of Agriculture, Forestry and Natural Environment, Aristotle University of Thessaloniki, P.O. Box261, Thessaloniki GR-54124, Greece

³Department of Electrical and Computer Engineering, Aristotle University of Thessaloniki, Thessaloniki 54 124, Greece

⁴Centre for Clinical, Experimental Surgery & Translational Research, Biomedical Research Foundation Academy of Athens, Athens, Greece

⁵Department of Biology, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand

⁶Institute of Plant Breeding and Genetic Resources ELGO-DEMETER (ex NAGREF), Thessaloniki, Macedonia, GR-57001, Greece

[#]Corresponding authors, the authors declare equal authorship.

email: nianiou@agro.aouth.gr

pmadesis@certh.gr

Download English Version:

<https://daneshyari.com/en/article/5589227>

Download Persian Version:

<https://daneshyari.com/article/5589227>

[Daneshyari.com](https://daneshyari.com)