Accepted Manuscript

miR-199a-5p and miR-495 target GRP78 within UPR pathway of lung cancer

Azam Ahmadi, Behzad Khansarinejad, Saman Hosseinkhani, Mostafa Ghanei, Seyed Javad Mowla

PII: S0378-1119(17)30211-1

DOI: doi: 10.1016/j.gene.2017.03.032

Reference: GENE 41838

To appear in: Gene

Received date: 9 December 2016 Revised date: 22 March 2017 Accepted date: 24 March 2017

Please cite this article as: Azam Ahmadi, Behzad Khansarinejad, Saman Hosseinkhani, Mostafa Ghanei, Seyed Javad Mowla , miR-199a-5p and miR-495 target GRP78 within UPR pathway of lung cancer. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Gene(2017), doi: 10.1016/j.gene.2017.03.032

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

miR-199a-5p and miR-495 target GRP78 within UPR pathway of lung cancer

Azam Ahmadi^a, Behzad Khansarinejad^b, Saman Hosseinkhani^c, Mostafa Ghanei^d, Seyed Javad Mowla^a,*

Running head: Hsa-miR-199a-5p and hsa-miR-495 target GRP78 in UPR pathway

Grant sponsor: This study supported by a PhD grant to the first author from Tarbiat Modares University.

Competing interests: The authors declare no conflicts of interest.

*Corresponding author: Seyed Javad Mowla, Professor, Molecular Genetics Department, School of Biological Sciences, Tarbiat Modares University, P.O. Box. 14115/175, Tehran, Iran, Tel.: (+98-21) 82883464, E-mail: sjmowla@modares.ac.ir

High lights:

- miRNAs targeting GRP78 within UPR pathway is proposed.
- GRP78 is up-regulated in tissue lung tumors, compared to non-tumor samples. miR-199a-5p and miR-495 are down-regulated in lung tumor samples compared to nontumor samples.
- Down-regulation of these miRNAs, might be the reason for up-regulation of GRP78 in NSCLC patients.
- miR-199a-5p and miR-495 have interaction with 3'-UTR of GRP78.
- These miRNAs might have effect on ER stress in lung cancer, and can be used as diagnostic biomarkers for activated UPR in NSCLC.

^a Department of Molecular Genetics, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran

^b Infectious Diseases Research Center (IDRC), Department of Microbiology and Immunology, Arak University of Medical Sciences, Arak, Iran

^c Department of Biochemistry, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, Iran

^d Research center of chemical injuries, Baqiyatallah University of Medical Sciences, Tehran, Iran

Download English Version:

https://daneshyari.com/en/article/5589524

Download Persian Version:

https://daneshyari.com/article/5589524

<u>Daneshyari.com</u>