

## Review

## Role of neuropilin-2 in the immune system

S. Schellenburg<sup>a</sup>, A. Schulz<sup>a</sup>, D.M. Poitz<sup>b</sup>, M.H. Muders<sup>a,\*</sup><sup>a</sup> Institute of Pathology, University Hospital Carl Gustav Carus, University of Technology, Dresden, Germany<sup>b</sup> Department of Internal Medicine and Cardiology, University Hospital Carl Gustav Carus, University of Technology, Dresden, Germany

## ARTICLE INFO

## Keywords:

Neuropilin-2

Polysialylation

## ABSTRACT

Neuropilins (NRPs) are single transmembrane receptors with short cytoplasmic tails and are dependent on receptors like VEGF receptors or Plexins for signal transduction. NRPs are known to be important in angiogenesis, lymphangiogenesis, and axon guidance. The Neuropilin-family consists of two members, Neuropilin-1 (NRP1) and Neuropilin-2 (NRP2). They are up to 44 % homologous and conserved in all vertebrates. High levels of NRP2 are found on immune cells. Current research is very limited regarding the functions of NRP2 on these cells. Recent evidence suggests that NRP2 is important for migration, antigen presentation, phagocytosis and cell-cell contact within the immune system. Additionally, posttranslational NRP2 modifications like polysialylation are crucial for the function of some immune cells. This review is an overview about expression and functions of NRP2 in the immune system.

## 1. Molecular structure of neuropilin-2

Neuropilins (NRPs) were first described in the optic nerve fibers of *Xenopus laevis* and named after the specific area where the antibody A5 binds, called “neuropile” (Takagi et al., 1987). There are two proteins representing the NRP family, NRP1 and NRP2. In humans, they are located on chromosome 10p11.22 for NRP1 and on chromosome 2q33.3 for NRP2. The NRPs are single transmembrane receptors with a predominant extracellular region containing two CUB domains (a1/a2), two Factor V/VIII homology domains (b1/b2), and a MAM domain (c) (Fig. 1). NRP1 and NRP2 can form homodimers as well as heterodimers. Both NRPs are highly conserved in vertebrates and the sequence homology of human and murine NRP2 is 94%. NRP1 and NRP2 are up to 44% homologous (Rossignol et al., 2000). Both molecules are heavily glycosylated. Additionally, NRP2 is one of the few proteins which can be sialylated (Curreli et al., 2007). The NRP1 and NRP2 have different splice variants which are between 551 and 926 amino acids in length. Two major variants exist in the case of NRP2, and are categorized as NRP2a and NRP2b. These differ in their intracellular C-terminal part (Fig. 1). In detail, the short C-terminally located cytoplasmic tail of NRP2a consists of 42 amino acids and has a PDZ-binding-domain with the C-terminal SEA amino acid sequence. The C-terminus of the other isoform, NRP2b, consists of 46 amino acids and shares just about 11% of the intracellular and transmembrane sequence of NRP2a (Fig. 1).

In mice, NRP2a can have insertions after amino acid 809 and the resulting variants are named according to the number of added amino acids, NRP2a (0), (5), (17) and (22). NRP2a (17) is generated by


alternative splicing between exon 15 and 16. For NRP2a (22) a different splice acceptor is active at intron 15 and 5 amino acids (GENFK) are added without shifting the reading frame (Fig. 1). Four splice variants for NRP2a were described but Rossignol et al. found only two isoform bands in lung, brain, heart, kidney, liver, placenta, and trachea in their PCR for NRP2a corresponding to NRP2a (17) and NRP2a (22). They weren't able to identify NRP2a (0) and NRP2a (5) in humans, which would be generated by splicing at exon 16, but the theoretical sequences can be found on PubMed. The theoretical human sequence for NRP2a (0) was used in a study from Rollenhagen et al., although it has never been identified in human tissue and cells at this point (Rollenhagen et al., 2013; Rossignol et al., 2000). It is possible that NRP2a (0) and NRP2a (5) are expressed in certain tissue or cells, but a more detailed analysis needs to be performed to shed light on the expression pattern of the splice variants of NRP2.

Rossignol et al. identified an exon sequence in intron 15 named exon 16b. It consists of 912 bp in total with a coding sequence of 283 bp, a stop codon and a 3'UTR including a cleavage and polyadenylation signal. Therefore, the alternative splicing for NRP2b results most likely between exon 15 and exon 16b. Two splice variants can occur, NRP2b (0) and NRP2b (5) depending on the insertion of 0 or 5 amino acids (GENFK) after amino acid 808 as described for NRP2a (Fig. 1) (Rossignol et al., 2000).

The difference in the C-terminal sequence, more specifically the PDZ domain, between the splice variants might result in different binding capabilities to other proteins. One protein with repetitive PDZ domains is GAIP interacting protein 1 (GIPC1)/Synectin. In fact, GIPC1/Synectin

\* Corresponding author.

E-mail address: [muders.michael@icloud.com](mailto:muders.michael@icloud.com) (M.H. Muders).



**Fig. 1.** Splice variants of Neuropilin 2.

Neuropilin 2 consists of two CUB domains followed by two Factor V/VIII domains followed by a MAM (c) domain for both NRP2a and NRP2b. These two are the main isoforms in human and mice. Between the MAM domain and the membrane (tm) additional splicing can occur and 5 additional amino acids (GENFK) can be added to NRP2a and NRP2b resulting in two different variants for NRP2a (17), (22) and two different transmembrane variants for NRP2b (0), (5) as well as one soluble form called sNRP2b. For mice, the following variants have been identified: NRP2a (0), (5), (17), (22) and NRP2b (0), (5).

binds to the C-terminal SEA domain and could be an initial trigger for a direct signal (Muders et al., 2006, 2009; Prahs et al., 2008). Furthermore, GIPC1/Synectin might have an important function as a scaffold for multiprotein complexes. Others and we have already shown that GIPC1/Synectin has important biological functions (Gao et al., 2000; Muders et al., 2009; Naccache et al., 2006). NRP2a is preferentially expressed in the brain, liver, lung, small intestine, kidney and heart while NRP2b is mainly expressed in the heart and muscles (Rossignol et al., 2000).

## 2. Binding and interaction partners

The group of soluble semaphorin 3 (SEMA 3) proteins and vascular endothelial growth factors (VEGFs) are known binding partners for the NRP family. NRP1 and NRP2 have different affinities to bind various semaphorins or VEGFs. Especially, SEMA3A binds to NRP1 whereas SEMA3F and SEMA3C bind to NRP2. VEGF-A binds to NRP1 and NRP2 but the affinity depends on the splice variants of VEGF-A. VEGF-C is an important binding partner for NRP2 (reviewed in (Prud'homme and Glinka, 2012)).

Semaphorins and VEGF have different binding domains and therefore are no competitors for binding to NRPs (Appleton et al., 2007). Also, it is well established that SEMA3 and VEGF binding to NRPs leads to dissimilar effects in the cell suggesting that different signal cascades become activated through binding of each ligand. SEMA3 are involved in vascular permeability, inhibiting of endothelial cell proliferation, and can induce apoptosis, whereas VEGFRs are involved in vasculogenesis and angiogenesis.

Another interaction partner for NRP2 is PTEN. PTEN is important for the inhibiting effect of the Semaphorin3F/NRP2 axis in some cells like T lymphocytes. In these cells, PTEN knock-down eliminates the inhibiting effect of Semaphorin3F (Nakayama et al., 2015).

A more detailed overview about the binding partners of NRP1 and NRP2 were reviewed 2012 from Prud'homme and Glinka (Prud'homme and Glinka, 2012).

NRPs are known for their crucial function in neurogenesis and cardiovascular development as it was shown in different animal models. For example, a full knock-out of NRP1 leads to death at embryonic day E10–E13.5 with a spectrum of cardiovascular defects (Takashima et al., 2002b). A knock-out of NRP2 is not lethal but displays abnormal guidance of nerves and fewer small vessels and capillaries. Double knock-out of NRP1 and NRP2 leads to an even earlier death at E8 and has a similar phenotype like a VEGF-A and VEGFR2 knock-out and therefore underlines the importance of the interactions between VEGFs and NRPs (Takashima et al., 2002a).

## 3. Posttranscriptional modification of NRP2 by ST8Sia IV

NRP2 and to a much lesser extent NRP1 are two of the few proteins which can be posttranslationally modified by polysialylation (Bhide et al., 2016; Mühlenhoff et al., 1998). To our knowledge only six other proteins in mammals can carry polysialylation. The neuronal adhesion molecule (NCAM/CD56) was first described in 1982 to carry polysialylations. Many researchers studied it during the years and were able to point out the importance of polysialic acid (polySia) (Finne, 1982) (reviewed in (C. E. Galuska et al., 2017)). Other polysialylated proteins are: the cluster of differentiation (CD) 36 (mouse/human) (Yabe et al., 2003), C-C chemokine receptor type 7 (CCR7) (mouse/human) (Kiermaier et al., 2016), E-selectin ligand-1 (ESL-1) (mouse/human) (Werneburg et al., 2016), Synaptic cell adhesion molecule SynCam-1 (mouse) (S. P. Galuska et al., 2010) and the sodium channel  $\alpha$  subunit (rat) (Zuber et al., 1992).

The CMP-N-acetylneuraminase-poly-alpha-2,8-sialyltransferase (ST8Sia IV) is one of two enzymes facilitating sialylation and is uniquely responsible for the addition of the sialic acid to NRP2. The other enzyme is called alpha-2,8-sialyltransferase 8B (ST8Sia II) and is not involved in the modification of NRP2 (Curreli et al., 2007). Addition of sialic acids can occur in a N-linked or O-linked manner. In case of NRP2, the sialylation is restricted to the linker region and can contain up to 100  $\alpha$ 2,8-linked sialic acid residues but has at least 8 residues (Rollenhagen et al., 2013; Sato and Kitajima, 2013). Despite 3 possible N-glycosylation sites and 2 N-glycans at Asn-152 and Asn-157, NRP2 carries only O-linked polysialylation (Rollenhagen et al., 2013). In the linker region, 6 possible sites for O-glycosylation and sialylation exist but the major, if not exclusive, sites are Thr 613, Thr 614, Thr 615 and Thr 619 (Rollenhagen et al., 2013) (Fig. 2). The sialylation varies between the different splice forms of NRP2. NRP2b and especially NRP2b (5) is more sialylated than NRP2a (Rollenhagen et al., 2013). Therefore, Bhide et al. elucidated the responsible domain for sialylation. They found that a replacement of the MAM domain of NRP2 by the MAM domain of NRP1 reduced the polysialylation of NRP2 significantly. Furthermore, they identified the Glu<sup>652</sup> and Glu<sup>653</sup> in the MAM domain as key elements for sialylation by ST8SiaIV (Bhide et al., 2016).

The polySia can bind to chemokines like CCL21 and the receptor CCR7 and is involved in migration. Other functions of the polySia are anti-adhesion, development of the nervous system, cell-cell and cell-matrix interaction, phagocytosis, and T-cell activation (Bhide et al., 2016; Drake et al., 2009; Rutishauser, 2008; Stamatatos et al., 2014).

In this review, we focus on the expression and known function of NRP2 in immune cells.

Download English Version:

<https://daneshyari.com/en/article/5591875>

Download Persian Version:

<https://daneshyari.com/article/5591875>

[Daneshyari.com](https://daneshyari.com)