



No evidence of an immune adjustment in response to a parasitoid threat in *Lobesia botrana* larvae



Fanny Vogelweith^{a,*}, Yannick Moret^b, Denis Thiéry^c, Lionel Delbac^c, Jérôme Moreau^b

^a Johannes-Gutenberg University of Mainz, Institute of Organismic and Molecular Evolution, Behavioral Ecology and Social Evolution Group, Johannes-von-Müller-Weg 6, 55128 Mainz, Germany

^b Université Bourgogne-Franche Comté, Équipe Écologie Évolutive, UMR 6282 Biogéosciences, 6 Bd Gabriel, 21000 Dijon, France

^c Institut National de la Recherche Agronomique (INRA), UMR 1065 Santé et Agroécologie du Vignoble (SAVE), Institut des Sciences de la Vigne et du Vin (ISVV), 71 avenue Edouard Bourlaux, 33883 Villenave d'Ornon Cedex, France

ARTICLE INFO

Keywords:

Campoplex capitator
Insect immunity
Grapevine moth
Phytomyia nigrina
Plasticity
Prophylaxis

ABSTRACT

Immune function is a key determinant of an organism's fitness, and natural insect populations are highly variable for this trait, mainly due to environmental heterogeneity and pathogen diversity. We previously reported a positive correlation between infection prevalence by parasitoids and host immunity in natural populations of the vineyard pest *Lobesia botrana*. Here, we tested whether this correlation reflects a plastic adjustment of host immunity in response to the local presence of parasites. To this end, we measured immunity of non-parasitized *L. botrana* larvae exposed, respectively, to one of the two most common species of parasitoids in vineyards, over 6 days. Larvae were able to sense the parasitoid through visual, chemical, or mechanical cues, but contact larvae-parasitoid were excluded. Contrary to our hypothesis, we found that *L. botrana* larvae did not increase their immune defenses in the presence of parasitoids, despite their ability to sense a potential threat. Our results therefore suggest that the positive correlation between infection prevalence by parasitoids and *L. botrana* immunity among natural populations may result from micro-evolutionary changes resulting from long-term local selection pressures imposed by parasitoids in wild populations rather than plastic adjustments of immunity.

1. Introduction

Animals live in dynamic environments and face variation in resource availability, climate and risk of infection over their whole lifespan. Thus, to survive and reproduce successfully, these organisms must allocate resources among competing physiological systems, such as immunity and growth (van der Most et al., 2011), to maximize fitness in changing environments. Immunity is one of the major physiological mechanisms regulating host survival (Lochmiller and Deerenberg, 2000). In insects, an important part of this defense relies on non-specific and constitutive mechanisms that involve the coordinated action of hemocytes and the phenoloxidase (PO) system (Siva-Jothy et al., 2005). Hemocytes are immune cells circulating in the hemolymph involved in the recognition and encapsulation of pathogens (Lavine and Strand, 2002). Conversely, PO mostly mediates the melanization of foreign objects and operates through the activation of the prophenoloxidase (PPO) cascade, its inactive precursor typically stored in the hemolymph and the hemocytes (Cerenius and Söderhäll, 2004).

Insect immunity is associated with inherent costs because it requires energy to build up, maintain and use (Armitage et al., 2003), and thus

reduces an individual's ability to invest into other physiological systems. Moreover, the activation of insect immunity produces chemical substances that are harmful for the producer and induce cumulative damage in its body (González-Santoyo and Córdoba-Aguilar, 2012; Nappi and Ottaviani, 2000). For this reason, individuals are expected to flexibly adjust their investment into the immune system to find an optimized balance between their ability to fight off pathogens on the one hand, and saving energy as well as limiting the accumulation of toxic immune components in their body on the other hand. Organisms can indeed be exposed to different threats such as parasites and parasitoids, which can vary among and between populations, as reported in *Drosophila melanogaster* (Kraaijeveld and Alphen, 1995; Tinsley et al., 2006; Corby-Harris and Promislow, 2008), *Gammarus pulex* (Cornet et al., 2009) and *Lobesia botrana* (Moreau et al., 2010; Vogelweith et al., 2013a). These different threats induce substantial variation in the investment in immunity within and among wild populations (Corby-Harris and Promislow, 2008; Cornet et al., 2009; Vogelweith et al., 2013a). For example, recent investigations among natural populations of the grapevine moth (*L. botrana*) revealed that levels of immune defenses in larvae were positively correlated to parasitoid infection

* Corresponding author.

E-mail address: fanny.vogelweith@gmail.com (F. Vogelweith).

prevalence (Vogelweith et al., 2013a). At a geographical scale, populations with high levels of immune defenses in non-parasitized larvae (high hemocyte concentrations, PPO enzyme activities and antimicrobial peptides) were also those exhibiting the highest rates of infection by parasitoids (Vogelweith et al., 2013a).

Although this confirms that parasitoid pressure strongly affects host immune investment (Bryan-Walker et al., 2007; Corby-Harris and Promislow, 2008), the exact cause of such a positive relationship remains unknown. Two non-mutually exclusive selective pressures could shape and drive such a variation in immune traits. First, parasites can mediate a local selection response in favor of (constitutively expressed) high levels of immune defense among hosts, which would result in positive covariation between parasite prevalence and immune defense (Lindström et al., 2004; Kalbe and Kurtz, 2006; Tschirren and Richner, 2006). Alternatively, such a positive correlation could arise if hosts plastically adjust their immune response to the (locally varying) risk of infection, since the ability to sense the current risk of parasitism through visual, chemical, or mechanical cues has been reported in some insect species (Peacor, 2003; Fievet et al., 2008). For example, when the risk of parasitism increases, insects show prophylactic investment in their immune defense even in the absence of parasites (Barnes and Siva-Jothy, 2000; Wilson and Reeson, 1998). The velvetbean caterpillar (*Anticarsia gemmatilis*) increased investment in primary defense barriers against parasites, such as the midgut, in response to increased conspecific density and an increased risk of infection (Silva et al., 2016). Such a plastic modulation of immune defenses can be rapid as reported in adult bumble-bee workers (*Bombus terrestris*) (Ruiz-González et al., 2009). A plastic investment in immunity is then likely to result in positive covariation between parasite prevalence and immune defense. To understand how parasitoids can modulate insect immunity, it is important to know which of the two above detailed selective pressures shapes investment into insect immunity.

To our knowledge, no study experimentally investigated how insect can plastically modulate their immune defenses in response to parasitoid pressure. Here, we test the hypothesis whether the presence of parasitoids in the local environment induces a plastic increase of the investment in immune defense. To this end, we exposed *L. botrana* larvae to either parasitoids (excluded physical contacts), or no parasitoid for a period of six days and measured immune parameters (hemocyte concentration and PPO activities system) on the last day of exposure. If *L. botrana* larvae indeed flexibly adjust their immunity to the parasitoid threat, we expected an increase of all immune parameters in the presence of parasitoids. In this scenario, the larvae of *L. botrana* would assess cues related to the presence of parasitoids and adjust their investment in immunity to match any increased threat of infection. *L. botrana* appears to be a good candidate because (1) a variation of larval immune parameters with the parasitoid pressure has already been reported among wild populations (Vogelweith et al., 2013a) and (2) the ability of *L. botrana* larvae to sense the current in the risk of parasitism has been previously demonstrated (Vogelweith et al., 2013b).

2. Material and methods

2.1. Model insect

The European grapevine moth *L. botrana* (Lepidoptera: Tortricidae) is currently the major grape pest in Europe and has also spread to American viticulture (Gilligan et al., 2011). This grapevine moth is a polyphagous insect that completes 2–5 generations a year (depending on latitude) on different grape cultivars, where the larvae can do considerable damage (Thiéry, 2008). The larvae used for this experiment come from inbred stock maintained diapause-free at the French National Institute for Agricultural Research (INRA) (Villenave d'Ornon, France) for more than 10 years. This strain is based on a great number of caged adults (several thousand per week) to which wild adults are periodically added. This laboratory strain has conserved some plasticity

because considerable variation is found in immune parameters between larvae (Muller et al., 2015). In addition, a very similar pattern in terms of basal immunity, response to an immune challenge and parasitoid escape behavior have been found between inbred stock and wild populations sampled in French vineyards (Vogelweith et al., 2014). We used larvae from the laboratory to ensure that they were not previously exposed to parasitoids because this could have affected their immunity. Larvae were fed with an ad libitum amount of a semi-artificial diet (Vogelweith et al., 2015), and maintained in boxes (18 × 11.5 × 7 cm) under standard laboratory conditions (22 ± 1 °C, 70 ± 10% r.h., and a L16:D8 photoperiod) at a density of approximately 100 individuals per 300 ml of diet.

2.2. Parasitoid populations

To mimic imminent parasitoid attack on the larvae, we used the two most common parasitoids of *L. botrana* larvae: *Campoplex capitator* (Hymenoptera: Ichneumonidae) and *Phytomyptera nigrina* (Diptera: Tachinidae) (Moreau et al., 2010). These parasitoids are known to be larval-parasitoid only and parasitize larvae of *L. botrana* from the 2nd to the 4th instar in vineyards in the south of France where they occur in sympatry (Thiéry, 2008). The parasitism rate of these two parasitoids strongly depends on the year and the vineyard considered (Moreau et al., 2010; Vogelweith et al., 2013a). For instance, in 2012, we observed a parasitism rate of 10% by *C. capitator* and 5% by *P. nigrina* while in 2011, we found a parasitism rate of 90% by *C. capitator* in the same vineyard (Vogelweith, unpublished data). Both parasitoid species were obtained from parasitized *L. botrana* larvae collected on Grenache in a vineyard close to Perpignan (southern France) in May 2012 using the procedure by Vogelweith et al. (2013a). In brief, the larvae were collected at the end of their development in grapes and kept in large polyethylene boxes (60 × 40 × 21.4 cm) in the laboratory where they were checked daily for pupation. Pupae were placed individually in glass tubes (70 × 9 mm diameter) closed with cotton plugs and stored under standard laboratory conditions (22 ± 1 °C, 70 ± 10% r.h., and L16:D8 photoperiod). The tubes were checked daily for the emergence of either the moth or the parasitoid. All the parasitoids were of the same age and were used two days after their emergence in the experiment.

2.3. Experimental design

Fourth instar larvae from the laboratory culture (see above) were exposed in groups of five to one female parasitoid (either *C. capitator* or *P. nigrina*) to assess the effect of parasitoid presence on larval immune parameters. In a previous experiment using then same protocol (Vogelweith et al., 2013b), we did not find any difference between control groups either exposed to a non-parasitoid fly or nothing. For this reason, we used additional groups of unexposed *L. botrana* larvae (empty cup) as sole control. The groups of larvae were placed in plastic boxes (98 × 98 × 49 mm) containing 50 ml of semi-artificial diet (Vogelweith et al., 2015) (Fig. 1). A plastic cup (30 mm diameter and 30 mm height) containing one female of *C. capitator* or *P. nigrina* (n = 6 parasitoids and n = 30 larvae, respectively), or no parasitoid (n = 5 and n = 20 larvae) was positioned in the middle of each box. The central cup was transparent and pierced with small holes, which allowed for visual, olfactory and vibratory stimulation of *L. botrana* larvae by the parasitoid. The parasitoids were offered a drop of honey as food for the duration of the experiment. The experimental boxes were maintained in standard laboratory conditions (22 ± 1 °C, 70 ± 10% r.h., and L16:D8 photoperiod). Six days later, a sample of hemolymph was collected from each larva for measurements of the hemocyte concentration and the activity of the PO-PPO system as described in Vogelweith et al. (2011). The larvae that reached metamorphosis (5%) did not provide hemolymph and were equally distributed among treatments. Note that we did not directly expose the larvae to the parasitoid because of the very aggressive behavior of *C. capitator*; since

Download English Version:

<https://daneshyari.com/en/article/5593089>

Download Persian Version:

<https://daneshyari.com/article/5593089>

[Daneshyari.com](https://daneshyari.com)