Accepted Manuscript

Coconut oil has less satiating properties than medium chain triglyceride oil



R. Kinsella, T. Maher, M.E. Clegg

PII:	S0031-9384(17)30211-1
DOI:	doi: 10.1016/j.physbeh.2017.07.007
Reference:	PHB 11849
To appear in:	Physiology & Behavior
Received date:	5 February 2017
Revised date:	25 June 2017
Accepted date:	5 July 2017

Please cite this article as: R. Kinsella, T. Maher, M.E. Clegg, Coconut oil has less satiating properties than medium chain triglyceride oil. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Phb(2017), doi: 10.1016/j.physbeh.2017.07.007

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Coconut oil has less satiating properties than medium chain triglyceride oil

R. Kinsella, T. Maher and M.E. Clegg*

Functional Food Centre, Department of Sport and Health Sciences, Faculty of Health and Life Sciences, Oxford Brookes University, **Potential mechanisms** Gipsy Lane, Oxford OX3 0BP, UK

*Corresponding author: Miriam Clegg, Functional Food Centre, Department of Sport and

Health Sciences, Faculty of Health and Life Sciences, Oxford Brookes University, Gipsy Lane,

Oxford OX3 0BP, UK

Email: mclegg@brookes.ac.uk; Ph: +44 1865 484365

Running title: MCT, coconut oil and satiety

Abstract

Download English Version:

https://daneshyari.com/en/article/5593671

Download Persian Version:

https://daneshyari.com/article/5593671

Daneshyari.com