


# Accepted Manuscript

Point: “Should acute fluid resuscitation be guided primarily by inferior vena caval ultrasound for patients in shock?” Yes

Gregory A. Schmidt, MD, FCCP



PII: S0012-3692(16)62457-1

DOI: [10.1016/j.chest.2016.11.021](https://doi.org/10.1016/j.chest.2016.11.021)

Reference: CHEST 829

To appear in: *CHEST*

Received Date: 8 November 2016

Accepted Date: 8 November 2016

Please cite this article as: Schmidt GA, Point: “Should acute fluid resuscitation be guided primarily by inferior vena caval ultrasound for patients in shock?” Yes, *CHEST* (2016), doi: 10.1016/j.chest.2016.11.021.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Point: “Should acute fluid resuscitation be guided primarily by inferior vena caval ultrasound for patients in shock?” Yes

Gregory A. Schmidt, MD, FCCP  
Division of Pulmonary Diseases, Critical Care, and Occupational Medicine  
Department of Internal Medicine  
University of Iowa, Iowa City, IA  
(319)384-6746

[Gregory-a-schmidt@uiowa.edu](mailto:Gregory-a-schmidt@uiowa.edu)

No conflicts of interest to report

Download English Version:

<https://daneshyari.com/en/article/5600989>

Download Persian Version:

<https://daneshyari.com/article/5600989>

[Daneshyari.com](https://daneshyari.com)