




Contents lists available at ScienceDirect

International Journal of Cardiology

journal homepage: www.elsevier.com/locate/ijcard

Cognitive ability, lifestyle risk factors, and two-year survival in first myocardial infarction men: A Swedish National Registry study[☆]

John Wallert^{a,*,1}, Guy Madison^{b,1}, Claes Held^{c,1}, Erik Olsson^{a,1}

^a Department of Public Health and Caring Sciences, Uppsala University, Uppsala, Sweden

^b Department of Psychology, Umeå University, Umeå, Sweden

^c Uppsala Clinical Research Center, Uppsala University, Uppsala, Sweden/Department of Medical Sciences: Cardiology, Uppsala University, Uppsala, Sweden

ARTICLE INFO

Article history:

Received 28 September 2016

Received in revised form 30 November 2016

Accepted 20 December 2016

Available online xxx

Keywords:

Behaviour and behavioural mechanisms

Cardiovascular disease

Intelligence

Lifestyle

Risk factors

Secondary prevention

ABSTRACT

Background: General cognitive ability (CA) is positively associated with later physical and mental health, health literacy, and longevity. We investigated whether CA estimated approximately 30 years earlier in young adulthood predicted lifestyle-related risk factors and two-year survival in first myocardial infarction (MI) male patients.

Methods: Young adulthood CA estimated through psychometric testing at age 18–20 years was obtained from the mandatory military conscript registry (INSARK) and linked to national quality registry SWEDEHEART/RIKS-HIA data on smoking, diabetes, hypertension, obesity (BMI > 30 kg/m²) in 60 years or younger Swedish males with first MI. Patients were followed up in the Cause of Death registry. The 5659 complete cases (deceased = 106, still alive = 5553) were descriptively compared. Crude and adjusted associations were modelled with logistic regression.

Results: After multivariable adjustment, one SD increase in CA was associated with a decreased odds ratio of being a current smoker (0.63 [0.59, 0.67], $P < 0.001$), previous smoker (0.79 [0.73, 0.84], $P < 0.001$), having diabetes (0.82 [0.74, 0.90], $P < 0.001$), being obese (0.90 [0.84, 0.95], $P < 0.001$) at hospital admission, and an increased odds ratio of two-year survival (1.26 [1.02, 1.54], $P < 0.001$). CA was not associated with hypertension at hospital admission (1.03 [0.97, 1.10], $P = 0.283$).

Conclusions: This study found substantial inverse associations between young adulthood CA, and middle-age lifestyle risk factors smoking, diabetes, and obesity, and two-year survival in first MI male patients. CA assessment might benefit risk stratification and possibly aid further tailoring of secondary preventive strategy.

© 2016 The Authors. Published by Elsevier Ireland Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Coronary Heart Disease (CHD) remains the leading cause of death globally and Myocardial Infarction (MI) its most common acute event, afflicting seven million individuals annually [1,2]. Acute MI and underlying CHD is a result of both genetic and lifestyle-related factors. The INTERHEART study found that nine modifiable risk factors, including smoking, unhealthy diet, diabetes, hypertension, physical inactivity, and psychological stress explained 90% of the population attributable risk of MI [3]. Patient behaviour is essential for reducing these modifiable

risk factors, and underlying causes of such behaviour is therefore of great importance.

General cognitive ability (CA) denotes the distillate of an individual's mental capacities [4–6]. CA is evident in basic day-to-day cognitive functions such as information processing, problem solving, and memory [7,8]. Within an individual, CA is highly stable from age 18 to 65 [9]. Quantified as the full-scale Intelligence Quotient (IQ), the intra-individual stability of CA displays the highest test-retest reliability of all psychological constructs [8,10]. Across individuals, however, CA exhibits considerable variation [8], which in turn is strongly predictive of several health-related life outcomes, for example educational achievement [11,12], job status attainment, and work performance [8,13,14]. Genetic studies have shown that CA is strongly heritable, and its phenotypic expression is subject to physical trauma and ageing effects, yet from adolescence and onwards it is resilient to other environmental influences (e.g. education) [8,15,16].

With respect to MI risk factors and mortality, aggregate research shows that higher CA early-in-life is associated with a greater likelihood to (i) exercise [17], (ii) eat healthy [17], (iii) be health literate [13], (iv)

[☆] Grant support: This work was supported by the Swedish Research Council for Health, Working Life, and Welfare [2014-4947], the Vårdal foundation [2014-0114], and U-CARE [2009-1093].

* Corresponding author at: Department of Public Health and Caring Sciences, Uppsala University, Box 564, Husargatan 3, SE 75122 Uppsala, Sweden.

E-mail address: john.wallert@pubcare.uu.se (J. Wallert).

¹ This author takes responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation.

be less prone to psychological stress and pathology [18], (v) not smoke [19], (vi) have less hypertension [20], (vii) not develop obesity [21], (viii) not develop CHD [22], and (ix) live longer [23]. This indicates that CA constitutes a general MI/mortality risk indicator, particularly since CA seems to predispose individuals for a range of cardiovascular risk factors.

It is therefore interesting to note that consensus guidelines on primary and secondary MI prevention [24,25] do not mention individual differences in CA, especially when there is evidence that current rehabilitation programs leave room for improvement [25,26]. In the SWEDEHEART national quality registry on Secondary Prevention after Heart Intensive Care (SEPHIA), only 21% of post-MI patients reach the four most important rehabilitation targets; smoking cessation, systolic blood pressure (SBP) < 140 mmHg, LDL cholesterol < 1.8 mmol/L, and maintaining adequate exercise [26]. Patient behaviour through lifestyle change is key in reaching these targets [25], and CA is in turn crucial for such target behaviour [13].

We related CA measured in young adulthood to four established lifestyle risk factors (smoking, diabetes, hypertension, and obesity) measured ~30 years later, and to two-year post-MI survival among first MI male patients ≤60 years. We hypothesized that higher CA was associated with: (1) lower prevalence of these four lifestyle risk factors at first MI hospital admission; (2) lower mortality within two years after the MI. We also hypothesized that (3) CA would be lower in the sample under study compared to simulated population data.

2. Methods

2.1. Data sources

The SWEDEHEART national quality Registry for Information and Knowledge about Swedish Heart Intensive Care Admissions (RIKS-HIA) contains data on >100 variables from patients admitted to a Cardiovascular Care Unit (CCU) in Sweden for acute coronary syndrome. RIKS-HIA is unselected and provides excellent coverage of the Swedish population (~90% of all MIs < 80 years). The local hospital decides on discharge diagnosis, for MI cases according to ICD codes I21–I23 [27] based on ECG, clinical symptoms, and other information [28]. The Swedish Cause of Death registry records the death date for patients that died during the study period. The Swedish National Archives (Riksarkivet) maintain the INSARK database which contains digitalized data from all Swedish men that performed mandatory military conscription at age 18–20 between years 1969–1997 [29]. Women were either not allowed or rarely conscripted, a limitation restricting our study to men [30]. Registries were linked by the National Board of Health and Welfare (Socialstyrelsen).

2.2. Ethical statement

These national registries aim to include all cases without discrimination, are sanctioned by Swedish law and monitored by designated registry organizations. All patients are informed of their inclusion and their right to have their data erased at any time at their own request. No one requested this during 2011 and 2013 and very few did before 2011. The present study was approved by the regional ethics committee in Uppsala (Dnr: 2013/478), and adheres to the 1975 Declaration of Helsinki.

2.3. Case selection and survival classification

We selected RIKS-HIA male patients that suffered a first MI in Sweden from 1st of January 2006 to 31st of December 2011. Each patient was classified as survivor or non-survivor depending on if the patient was alive 24 months after their MI admission date. Therefore, the maximal theoretical age of RIKS-HIA patients that also had INSARK data was 62 years (born 1949, conscript 1969 at age 20, and had a first MI in 2011 at age 62). Implementation delays of the conscript procedure and the scarcity of 20 year old conscripts rendered an effective sample ≤ 60 years of age at the time of hospital admission. Hence, CA was estimated when patients were between 18 and 20, lifestyle outcomes were measured at the time of hospital admission when these subjects were admitted for their first MI, i.e. 30–60 years old (mean age = 51.8), and mortality was assessed two years post-MI. The average age difference between the predictor (CA) and outcomes was therefore approximately 30 years. There were few incomplete cases with missing values, primarily for length (16.9%), weight (7.4%), SBP (5.9%), HR (4.2%), and smoking (3.1%). Removal of incomplete cases rendered a final sample of 5659 complete cases (106 deceased, 5553 still alive). For reference, we also calculated baseline characteristics for 34,849 cases constituting the whole first MI male population registered between 2006 and 2011 in RIKS-HIA with complete values on all but the CA variable.

2.4. Psychometric variables

We used the standardized nine (Stanine) scaled individual performance on the four psychometric tests (S1–4) from the Swedish Enlistment Battery (SEB). These tests estimate verbal ability (S1), logical reasoning (S2), spatial/non-verbal ability (S3), and technical understanding (S4) [30]. A principal component analysis revealed similar and substantial factor loadings from all subtests on the first principal component, taken to be the general CA factor (loadings range = −0.53 to −0.46). This first composite factor was the only one which eigenvalue was >1 and accounted for most of the variance in test scores (65.6%). Similar to [9], we therefore added the four standardized subtest scores and divided their sum by four to construct a unit weighted CA estimate. Because the psychometric scores on each subtest were standardized in relation to the total male conscript cohort with $\mu = 5$, $\sigma = 2$, and are normally distributed in this population [30], it was possible to simulate CA in the full conscript cohort. This required calculating the population standard deviation (σ) which was done by taking the ratio of the population mean (μ) and the sample mean (\bar{x}) and multiplying this with the sample standard deviation (s)

$$\sigma = \left(\frac{\mu}{\bar{x}}\right) * s$$

which resulted in $\sigma = 1.56$. We then simulated a Gaussian vector of length = 5659 with $\mu = 5$ and $\sigma = 1.56$ to represent CA in the full conscript cohort.

2.5. Lifestyle risk factors and covariates

We selected four established MI/mortality risk factors with clear behavioural components, also found to be CA-related in previous studies [5,19,21,31]: self-reported smoking (current smoker/previous i.e. quit > 1 month ago/never), known comorbid diabetes (yes/no), hypertension defined as if previously or presently medicated for this condition (yes/no), and obesity defined as body mass index (BMI) > 30 kg/m² (yes/no). Additionally, we selected age (years), resting-state SBP (mmHg), resting-state heart rate (bpm), primary percutaneous coronary intervention (PCI) or coronary artery bypass grafting (CABG) (yes/no), and discharge medicines: ACE-inhibitors and/or A2-blockers (yes/no), oral anti-coagulants (yes/no), other antiplatelet (yes/no), beta blockers (yes/no), and statins (yes/no).

2.6. Statistical procedures and software

We present continuous variables as mean (SD) and categorical variables as count (%) unless specified otherwise. For group comparisons on categorical variables we used the Pearson's χ^2 -test, and for continuous variables the Welch's *t*-test. Binomial and multinomial logit modelling was performed to estimate the adjusted influence of CA on the odds of lifestyle risk factor prevalence and survival. Odds ratios (OR) with 95% profile likelihood confidence intervals (CI) and *z*-tests for coefficient significance are reported. Given the patients relatively low age, we only assumed a log-linear association between age and survival. Statistical significance was set to 5% (two-tailed). Complete cases were analysed in R (version 3.2.3, R Development Core Team, Austria, Vienna) [32] using packages *AER*, *base*, *MASS*, *nnet*, *plyr*, *psych*, *stats*, and *rms*.

3. Results

Age ranges for complete cases at first MI admission in the full sample (30–60), the subsample deceased (38–60), subsample still alive (30–60), and the total first MI male RIKS-HIA population (11–99) are reported complementary to the mean (SD) ages represented in Table 1. Descriptive statistics in Table 1 shows that all four psychometric mean estimates and their aggregated mean CA were lower in non-survivors compared to survivors (comparison of CA means, $P = 0.003$). Non-survivors were also slightly older ($P < 0.001$), more likely to currently smoke, less likely to never have smoked ($P = 0.007$), more likely to have diabetes ($P < 0.001$) at hospital admission, and less likely to receive beta blockers ($P < 0.001$), statins ($P < 0.001$), and other antiplatelets ($P < 0.001$) at discharge. Heart rate was higher ($P < 0.001$) in non-survivors. Group comparisons for previous smoking, hypertension, and obesity prevalence, primary PCI and CABG, and for height, weight, SBP, and BMI were not statistically significant. CA in the full study sample was also lower compared to CA in the simulated full male conscript population ($P < 0.001$). Compared to the empirical norm, we see that our full sample represents a considerably younger portion of first MIs in Sweden ($P < 0.001$).

Logit modelling was then performed and odds ratios computed (OR). Table 2 shows that one SD increase in CA corresponds to an unadjusted (crude) decreased odds by 37% of being a current smoker, 20% of being a previous smoker, 18% of having diabetes, 13% of being obese,

Download English Version:

<https://daneshyari.com/en/article/5605121>

Download Persian Version:

<https://daneshyari.com/article/5605121>

[Daneshyari.com](https://daneshyari.com)