

# Accepted Manuscript

Beta-adrenergic Receptors are Critical for Weight Loss but not for other Metabolic Adaptations to the Consumption of a Ketogenic Diet in Male Mice

Nicholas Douris, Bhavna N. Desai, ffolliott M. Fisher, Theodore Cisu, Alan J. Fowler, Eleen Zarebidaki, Ngoc Ly T. Nguyen, Donald A. Morgan, Timothy J. Bartness, Kamal Rahmouni, Jeffrey S. Flier, Eleftheria Maratos-Flier

PII: S2212-8778(17)30307-1

DOI: [10.1016/j.molmet.2017.05.017](https://doi.org/10.1016/j.molmet.2017.05.017)

Reference: MOLMET 488

To appear in: *Molecular Metabolism*


Received Date: 28 April 2017

Revised Date: 26 May 2017

Accepted Date: 31 May 2017

Please cite this article as: Douris N, Desai BN, Fisher fM, Cisu T, Fowler AJ, Zarebidaki E, Nguyen NLT, Morgan DA, Bartness TJ, Rahmouni K, Flier JS, Maratos-Flier E, Beta-adrenergic Receptors are Critical for Weight Loss but not for other Metabolic Adaptations to the Consumption of a Ketogenic Diet in Male Mice, *Molecular Metabolism* (2017), doi: 10.1016/j.molmet.2017.05.017.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Beta-adrenergic Receptors are Critical for Weight Loss but not for other Metabolic Adaptations to the Consumption of a Ketogenic Diet in Male Mice**

Nicholas Douris<sup>\*1,4</sup>, Bhavna N. Desai<sup>\*1</sup>, ffolliott M. Fisher<sup>1</sup>, Theodore Cisu<sup>1</sup>, Alan J. Fowler<sup>1</sup>, Eleen Zarebidaki<sup>2</sup>, Ngoc Ly T. Nguyen<sup>2</sup>, Donald A. Morgan<sup>3</sup>, Timothy J. Bartness<sup>2</sup>, Kamal Rahmouni<sup>3</sup>, Jeffrey S. Flier<sup>1</sup>, Eleftheria Maratos-Flier<sup>1</sup>

\*Equally contributing authors

<sup>1</sup> Department of Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA 02215

<sup>2</sup> Department of Biology and Center for Obesity Reversal, Georgia State University, Atlanta, GA 30302-4010, USA

<sup>3</sup> Department of Pharmacology, University of Iowa, Carver College of Medicine, Iowa City IA 52242, USA

<sup>4</sup> Present address: Alkermes Inc., Waltham MA, USA

Correspondence to:

Eleftheria Maratos-Flier, MD  
Professor of Medicine  
Department of Medicine  
Beth Israel Deaconess Medical Center  
Harvard Medical School  
Center for Life Sciences  
3 Blackfan Circle  
Boston, MA 02215

Phone: 617-735-3289

Fax: 617-735-3343

[emaratos@bidmc.harvard.edu](mailto:emaratos@bidmc.harvard.edu)

Word count: 3850 (intro, method, results, discussion)

Figures: 8

Running title: SNS activation required for ketogenic diet induced weight loss

Download English Version:

<https://daneshyari.com/en/article/5618706>

Download Persian Version:

<https://daneshyari.com/article/5618706>

[Daneshyari.com](https://daneshyari.com)