



Review Article

Insulin resistance: vascular function and exercise

Moon-Hyon Hwang^{a,b}, Sewon Lee^{b,c,*}^a Division of Health and Exercise Science, Incheon National University, Incheon, Korea^b Sport Science Institute, Incheon National University, Incheon, Korea^c Division of Sport Science, Incheon National University, Incheon, Korea

ARTICLE INFO

Article history:

Received 12 May 2016

Received in revised form

30 May 2016

Accepted 2 June 2016

Available online 9 June 2016

Keywords:

aerobic exercise

atherosclerosis

diabetes mellitus

metabolic syndrome

resistance exercise

ABSTRACT

Insulin resistance associated with metabolic syndrome and Type 2 diabetes mellitus is an epidemic metabolic disorder, which increases the risk of cardiovascular complications. Impaired vascular endothelial function is an early marker for atherosclerosis, which causes cardiovascular complications. Both experimental and clinical studies indicate that endothelial dysfunction in vasculatures occurs with insulin resistance. The associated physiological mechanisms are not fully appreciated yet, however, it seems that augmented oxidative stress, a physiological imbalance between oxidants and antioxidants, in vascular cells is a possible mechanism involved in various vascular beds with insulin resistance and hyperglycemia. Regardless of the inclusion of resistance exercise, aerobic exercise seems to be beneficial for vascular endothelial function in both large conduit and small resistance vessels in both clinical and experimental studies with insulin resistance. In clinical cases, aerobic exercise over 8 weeks with higher intensity seems more beneficial than the cases with shorter duration and lower intensity. However, more studies are needed in the future to elucidate the physiological mechanisms by which vascular endothelial function is impaired in insulin resistance and improved with aerobic exercise.

© 2016 Korea Institute of Oriental Medicine. Published by Elsevier. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Insulin resistance is a metabolic disorder that reflects low capability to wash glucose out of the bloodstream to target tissues. Unlike type 1 diabetes mellitus, which results from disrupted insulin secretion from pancreatic β cells, type 2 diabetes mellitus is a representative metabolic disease attributable to increased insulin resistance and decreased insulin sensitivity.¹ Type 2 diabetes mellitus comprises

> 90% of all cases of diabetes and affected > 350 million people in 2011.² In particular, the population of diabetes mellitus is gradually increasing in Korea, and about 15,000 diabetic patients die every year owing to the associated complications.² Metabolic syndrome, a clinically prediabetic condition characterized by abdominal obesity, hypertension, hyperglycemia, and dyslipidemia, is closely associated with increased insulin resistance in etiology. Approximately 25% of the adult population is considered to be at risk of metabolic syndrome globally,³ and about 35% of the adult population in

* Corresponding author. Division of Sport Science and Sport Science Institute, Incheon National University, Building Number 16, Room Number 423, (Songdo-dong) 119 Academy-ro, Yeonsu-gu, Incheon, Korea.

E-mail address: leesew@inu.ac.kr (S. Lee).

<http://dx.doi.org/10.1016/j.imr.2016.06.001>

2213-4220/© 2016 Korea Institute of Oriental Medicine. Published by Elsevier. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

the United States is reported to have metabolic syndrome.⁴ In Korea, the population with metabolic syndrome increased from 25% in 1998 to 31% in 2007.⁵ Metabolic syndrome and type 2 diabetes mellitus are etiologically similar because they share two common causes: insulin resistance and abdominal obesity.⁶ Thus, patients with metabolic syndrome have about five times higher possibility to progress into type 2 diabetes mellitus compared with those without metabolic syndrome.⁷

Patients with type 2 diabetes mellitus or metabolic syndrome are usually at elevated risk of cardiovascular disease.^{2,8,9} Degenerative alterations in vascular structure and function in the metabolic diseases result in atherosclerosis, a main cause of cardiovascular disease.⁸ Atherosclerosis results from dysfunction of the vascular endothelium, which is closely related to decreased NO bioavailability and increased endothelium-derived contraction factors in blood vessels.¹⁰ Previous studies have indicated that excessive oxidative stress in the vasculature of patients with with metabolic disease is a possible mechanism to cause reduction in NO bioavailability and to impair vascular endothelial function.^{9,11–13} However, few studies have demonstrated the physiological mechanisms from vascular cells in patients with metabolic disease. Regular physical activity is known as a strategy to reverse vascular dysfunction and to prevent future cardiovascular disease. However, a few studies have investigated the effect of exercise training on vascular function in patients with type 2 diabetes mellitus and metabolic syndrome.^{14–17} Thus, the purpose of this review is to introduce potential physiological mechanisms by which vascular function is impaired in augmented insulin resistance, to summarize previous studies to investigate the effects of exercise training on vascular function in type 2 diabetes mellitus and metabolic syndrome, and to facilitate related studies in the future.

2. Insulin resistance and oxidative stress: a putative physiological mechanism to cause vascular dysfunction

Increased insulin resistance leads to augmented glucose level in the bloodstream. Since vascular endothelium is the single innermost layer in the vascular structure, vascular endothelial cells are likely to be damaged by hyperglycemic stress, suggesting that metabolic dysfunction such as insulin resistance and type 2 diabetic mellitus may result in vascular dysfunction.¹⁸ It has been demonstrated that the generation of reactive oxygen species (ROS) is increased in both large and small vascular beds collected from hyperglycemic animal models.¹⁹ In a hyperglycemic environment, elevated ROS generation in vascular cells can be influenced by various biological signaling pathways. Aldose reductase, in hyperglycemic conditions, converts glucose to sorbitol by using nicotinamide adenine dinucleotide phosphate as a cofactor, thus the regeneration of reduced glutathione is reduced and intracellular oxidative stress level is increased.^{2,20,21} Increased advanced glycation end-products lead to increased receptor activity, which facilitates ROS and inflammatory cytokine production.^{2,22,23} Activation of the diacylglycerol-protein kinase C pathway also contributes to the elevated generation of ROS and inflammatory cytokines.^{2,24–26} In the

overloaded glycolytic pathway induced by hyperglycemia, uridine diphosphate N-acetyl glucosamine has a negative effect on endothelial NO synthase activity by obstructing its phosphorylation at serine 1177 and increases the expression of proinflammatory cytokines such as transforming growth factor- β 1 and plasminogen activator inhibitor-1.^{2,27–29} In the vascular endothelium, oxidative stress induced by increased ROS generation not only directly plays a pivotal role in reducing NO bioavailability, but also leads to increased expression of proinflammatory cytokines so that proatherogenic and prothrombotic processes are abnormally facilitated.³⁰ Thus, the structural and functional integrity of vascular endothelium is likely to be damaged with insulin resistance, which increases the risk of cardiovascular disease in those who are continuously exposed to hyperglycemia in the circulation.

3. Insulin resistance and exercise: effects on vascular function

Insulin signaling in local vascular cells is essential to promote vascular endothelial NO production.³¹ Thus, the increase in insulin resistance in the vasculature has a negative effect on vascular endothelial function and cardiovascular morbidity and mortality. Regular physical activity and exercise are known to promote cardiovascular health in those with insulin resistance.

3.1. Effect of exercise training on vascular function in individuals with insulin resistance

There are no specific exercise guidelines for metabolic syndrome patients to preserve and promote their cardiovascular health. Even though their joint statements were published in both 2000 and 2010, the American College of Sport Medicine and the American Diabetes Association only provide generic exercise regimens for diabetic patients.^{32,33} In the two previous position standards, the experts recommended aerobic exercise or aerobic exercise combined with resistance exercise to prevent and promote cardiovascular disease risk in patients with diabetes. However, the effect of either aerobic or combined exercise training on both micro- and macrovascular endothelial function is still in debate. Some studies presented positive results of exercise training,^{17,34–36} but others did not show any positive effect of exercise on vascular health in type 2 diabetic patients.^{37–39} For individuals with metabolic syndrome, a small number of studies have explored the effect of exercise training on vascular endothelial function, but the training effect has presented a positive tendency,^{14,15,40,41} and had a close relationship with increased NO bioavailability.⁴² It is established that insulin resistance contributes to impaired vascular endothelial function,³¹ and either aerobic or combined exercise training programs might help to reverse endothelial dysfunction in those who have insulin resistance.^{14,15,17,34–36,40,41} Exercise intensity also seems to be a main factor influencing exercise effects in people with insulin resistance. Although both moderate-intensity continuous training and high-intensity interval training have shown improvement in vascular endothelial function, in the same caloric

Download English Version:

<https://daneshyari.com/en/article/5635141>

Download Persian Version:

<https://daneshyari.com/article/5635141>

[Daneshyari.com](https://daneshyari.com)