



Short Communication

Efficacy of herbal toothpastes on salivary pH and salivary glucose – A preliminary study



Mahesh R. Khairnar^{*1}, Arun S. Dodamani, G.N. Karibasappa², Rahul G. Naik, Manjiri A. Deshmukh³

Department of Public Health Dentistry, JMF's ACPM Dental College, Dhule, Maharashtra, India

ARTICLE INFO

Article history:

Received 30 August 2016

Received in revised form

11 October 2016

Accepted 22 December 2016

Available online 12 March 2017

Keywords:

Glucose

Toothpastes

Oral health

pH

Saliva

ABSTRACT

Due to dearth of literature on the effect of herbal toothpaste on saliva and salivary constituents, the present study was undertaken to evaluate and compare the effect of three different herbal toothpastes with the focus on on salivary pH and salivary glucose. Forty five subjects in the age group of 19–21 years were randomly divided into 3 groups (15 in each group) and were randomly intervened with three different herbal toothpastes (Dant Kanti, Himalaya Complete Care and Vicco Vajradanti). Unstimulated saliva samples were collected before and after brushing and salivary glucose and pH levels were assessed at an interval of one week each for a period of 4 weeks starting from day 1. All the three toothpastes were effective in reducing the overall ($p < 0.05$) levels as well as levels of salivary glucose from pre-brushing to post-brushing at each interval ($p < 0.05$) and in increasing the overall levels as well as levels of salivary pH ($p < 0.05$) from pre-brushing to post-brushing at each interval. Herbal toothpastes were effective in reducing salivary levels of glucose and improving pH of the saliva.

© 2016 Transdisciplinary University, Bangalore and World Ayurveda Foundation. Publishing Services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Mechanical oral hygiene procedures employing proper use of toothbrushes along with dentifrices seem to be the most effective form among all oral hygiene measures. Dentifrices serve as an ideal vehicle for any active therapeutic ingredient [1]. Among dentifrices, various types of toothpastes serve as the most common and accessible preventive oral health care tool available. It has been proven that modern toothpastes can help prevent dental caries and limit the regrowth of dental plaque & gingivitis [2].

Due to certain side-effects of routinely used non-herbal toothpastes such as mucosal ulceration, circumoral dermatitis, etc., people are resorting to herbal toothpaste which do not contain any synthetic ingredient [3,4]. Consumers who gravitate towards using herbal products often view these products as being safer than products that contain chemicals. Their efficacy can be attributed to various properties such as anti-inflammatory, anti-microbial, astringent action, anti-diabetic, anti-fungal, analgesic and antiseptic properties. However there is lack on professional consensus regarding use of herbal dental products due to lack of sufficient clinical research and literature on the efficacies of herbal based toothpastes.

Salivary components like pH, glucose which play an important role in maintaining oral health [5,6]. But scientific studies assessing the effect of non-herbal as well as herbal toothpastes on these salivary components are lacking. Most of the available literature has focused mainly on antimicrobial properties of herbal toothpastes [7]. The three toothpastes used in the study i.e., Patanjali Dant-Kanti Dental Cream, Himalaya Complete Care Toothpaste and Vicco Vajradanti Toothpaste were selected based on their popularity. Hence, the present study evaluates and compares the effect of these toothpastes on salivary pH and salivary glucose.

* Corresponding author. Department of Public Health Dentistry, Bharati Vidyapeeth Deemed University Dental College and Hospital, Sangli, Maharashtra, India.

E-mail address: kmahesh222@gmail.com (M.R. Khairnar).

Peer review under responsibility of Transdisciplinary University, Bangalore.

¹ Department of Public Health Dentistry, Bharati Vidyapeeth Deemed University Dental College and Hospital, Sangli, India.

² Department of Public Health Dentistry, Dr. DY Patil Dental School, Pune, India.

³ Department of Public Health Dentistry, Swargiya Dadasaheb Kalmegh Smruti Dental College & Hospital, Nagpur, India.

2. Material and methods

The present study was a double blind randomized study conducted on 45 BDS students (2nd and 3rd BDS students in the age group of 19–21 years old). Ethical clearance for the study was obtained from the Institutional Ethical Review Board and informed consent was obtained from all the participants prior to the study. Three herbal toothpastes tested were: Group A – Patanjali Dant Kanti Dental Cream; Group B – Himalaya Complete Care Toothpaste and Group C – Vicco Vajradanti Toothpaste.

2.1. Inclusion criteria

Participants in the age group of 19–21 years with good general health following routine oral hygiene procedures and agreement to comply with the study visits were included in the study.

2.2. Exclusion criteria

Participants suffering from systemic diseases, currently using antibiotics, undergoing orthodontic treatment, using additional oral hygiene aids other than toothbrush & toothpaste, having adverse habits like smoking, alcohol consumption were excluded from the study.

2.3. Study design

Since this was a pilot study, it was decided to include 15 participants in each group [8]. Forty-five participants from all the dental students who fulfilled inclusion and exclusion criteria were selected by simple random method and were randomly divided into three groups of 15 each. Three different toothpastes were assigned to three groups by simple random method.

Both, the participants and the investigator were blinded with respect to the study groups. All the study toothpastes were covered with white sticker label indicating only group name and caps of all the toothpastes were replaced with white caps of same uniform size to achieve blinding. The participants were instructed to brush for 3 min twice daily using the toothpaste which was provided to them. They were also asked not to use any other form of oral hygiene aids apart from toothpastes provided to them. Under the observation of the investigator, unstimulated saliva samples were collected in a container by the subjects before and immediately after brushing prior to rinsing with water and were transported to the lab where salivary pH and salivary glucose levels were determined. The study was conducted over a period of 4 weeks on each participant and saliva samples were obtained at weekly intervals starting from day 1 i.e. on 1st day, 7th day, 14th day, 21st day and 28th day.

Table 1
Mean change in salivary glucose (mg/dl).

Interval	Pre-brushing and post-brushing levels in salivary glucose					
	Dant kanti		Himalaya Complete Care		Vicco Vajradanti	
	Pre	Post	Pre	Post	Pre	Post
1st day	5.82 ± 0.4 P = 0.009	5.63 ± 0.5	5.89 ± 0.64 P < 0.001	5.60 ± 0.61	5.91 ± 0.41 P < 0.001	5.32 ± 0.37
7th day	5.77 ± 0.4 P < 0.001	5.42 ± 0.4	5.93 ± 0.72 P < 0.001	5.56 ± 0.60	5.82 ± 0.42 P < 0.001	5.24 ± 0.34
14th day	5.66 ± 0.4 P < 0.001	5.32 ± 0.4	5.83 ± 0.68 P < 0.001	5.52 ± 0.60	5.76 ± 0.41 P < 0.001	5.19 ± 0.35
21st day	5.57 ± 0.41 P < 0.001	5.24 ± 0.41	5.74 ± 0.73 P < 0.001	5.43 ± 0.66	5.68 ± 0.45 P < 0.001	5.09 ± 0.35
28th day	5.44 ± 0.42 P < 0.001	5.11 ± 0.39	5.62 ± 0.74 P < 0.001	5.26 ± 0.68	5.59 ± 0.44 P < 0.001	5.03 ± 0.37

2.4. Saliva collection and analysis

The required quantity of saliva sample was collected before and after tooth-brushing procedure. The participants were asked to refrain from eating and drinking for 1 h before sample collection. The unstimulated salivary samples (10 ml of saliva for each subject) were collected by the participants in sterile plastic container containing 0.1 ml of sodium fluoride in morning between 7 and 7.30 am before and after brushing by spitting method [9]. Each saliva sample was centrifuged at 3000 rpm for 10 min and clear supernatants were processed immediately for estimation of pH and glucose. Salivary pH was measured using a calibrated pH meter (Vanira Li 613 Digital pH meter) and salivary glucose levels were determined by Glucose oxidase/peroxidase reagent method using Erba Mannheim auto-analyzer as per standard protocol.

2.5. Statistical analysis

Data were analyzed using SPSS software version 16. Mean and standard deviation of the obtained pre (before brushing) and post (after brushing) levels of Salivary Glucose and Salivary pH at 1st, 7th, 14th, 21st and 28th day were compared within three groups by paired 't' test. Overall changes in the levels of salivary glucose and salivary pH on 1st and 28th day between three groups were compared by ANOVA test.

3. Results

3.1. Change in salivary glucose (mg/dl) and pH

All the three toothpastes showed significant change in salivary glucose and salivary pH levels after brushing at each interval (1st, 7th, 14th, 21st and 28th day) (Tables 1 and 2).

3.2. Overall difference in salivary glucose (mg/dl) and pH

Overall change in baseline (pre-brushing) values of salivary glucose from 1st day to 28th (last) day were found to be significant in all the three groups; however salivary pH values were found to be non-significant (Table 3).

4. Discussion

The present study compared the effects of three different herbal toothpastes on salivary pH and salivary glucose. The overall experience with the use of toothpaste reported was good with no adverse reactions. Reduction in salivary glucose levels and an increase in salivary pH levels were seen after use of allotted toothpastes. The present study incorporated only herbal toothpaste

Download English Version:

<https://daneshyari.com/en/article/5635277>

Download Persian Version:

<https://daneshyari.com/article/5635277>

[Daneshyari.com](https://daneshyari.com)