



## Green tea catechin inhibits the activity and neutrophil release of Matrix Metalloproteinase-9



Wan K. Kim-Park<sup>a</sup>, Eman S. Allam<sup>b, c</sup>, Jadesada Palasuk<sup>d, e</sup>, Michael Kowolik<sup>b</sup>,  
Kichuel K. Park<sup>f</sup>, L. Jack Windsor<sup>b, \*</sup>

<sup>a</sup> Department of Periodontology, Indiana University School of Dentistry, Indianapolis, IN, USA

<sup>b</sup> Department of Oral Biology, Indiana University School of Dentistry, Indianapolis, IN, USA

<sup>c</sup> Oral and Dental Research Division, National Research Centre, Cairo, Egypt

<sup>d</sup> Department of Restorative Dentistry, Division of Dental Biomaterials, Indiana University School of Dentistry, Indianapolis, IN, USA

<sup>e</sup> Department of Restorative Dentistry, Faculty of Dentistry, Naresuan University, Phitsanulok, Thailand

<sup>f</sup> Department of Preventive & Community Dentistry, Indiana University School of Dentistry, Indianapolis, IN, USA

### ARTICLE INFO

#### Article history:

Received 17 December 2014

Received in revised form

21 February 2015

Accepted 26 February 2015

Available online 29 April 2015

#### Keywords:

Matrix metalloproteinases

Periodontal diseases

Green tea catechin

Neutrophils

Epigallocatechin-gallate

### ABSTRACT

Green tea (*Camellia sinensis*; 綠茶 lǜ chá) extracts have been shown to possess anti-oxidant and anti-inflammatory effects in various cell types. Green tea extract (GTX) has been shown to significantly inhibit the activity of collagenase-3 (matrix metalloproteinase-13 (MMP-13)) *in vitro*. MMPs, such as MMP-9, are known to be involved in many inflammatory diseases including periodontal disease. GTX and a major catechin, epigallocatechin-gallate (EGCG), were examined for their ability to inhibit purified MMP-9 activity and its release from stimulated neutrophils.

Methanol extract of Green tea and commercially purchased EGCG (>95 % purity) were tested *in vitro* for their ability to inhibit MMP-9 activity and/or its release from neutrophils using a  $\beta$ -casein cleavage assay and gelatin zymography, respectively. Statistical analysis was performed by Student's t-test.

GTX and EGCG at 0.1% (w/v) completely inhibited the activity of MMP-9. In addition, GTX and EGCG (0.1 %) significantly inhibited ( $p < 0.001$ ) the release of MMP-9 from formyl-Met-Leu-Phe (FMLP)-stimulated human neutrophils by  $62.01\% \pm 6.717$  and  $79.63\% \pm 1.308$ , respectively. The inhibitory effects of GTX and EGCG occurred in unstimulated neutrophils ( $52.42\% \pm 3.443$  and  $62.33\% \pm 5.809$ , respectively). When the inhibitory effect of EGCG was further characterized, it significantly inhibited the release of MMP-9 from the FMLP-stimulated human neutrophils in a dose-dependent manner.

The effects of GTX and EGCG on MMPs could be extrapolated to clinical/*in vivo* studies for the development of oral care products to prevent or treat chronic inflammatory diseases including periodontal diseases.

Copyright © 2016, Center for Food and Biomolecules, National Taiwan University. Production and hosting by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### 1. Introduction

Bacterial invasion stimulates infiltrated neutrophils to produce reactive oxygen species (ROS). These ROS modulate various enzyme activities including protein kinases, ion channels, membrane receptors and transcriptional factors like nuclear factor kappa B

(NF $\kappa$ B). In turn, these stimulate the production of cytokines and host matrix metalloproteinases (MMPs).<sup>1</sup>

MMPs are a family of at least 27 zinc-containing endopeptidases that are classified as collagenases, gelatinases, stromelysins, membrane-type MMPs and others.<sup>2</sup> MMP-9 (gelatinase-B) is believed to be involved in many inflammatory diseases including periodontal diseases, tumor growth and metastasis, arthritis and cardiovascular diseases.<sup>3–6</sup> MMP-9 is expressed in neutrophils constitutively before they are released from the bone marrow to the circulation as part of the innate defense armory.<sup>7</sup> The expression of the MMPs also occurs at the tissue level after neutrophil migration upon infection in an inductive manner. MMP-9 is stored in neutrophils' granules and released upon activation.<sup>7</sup>

\* Corresponding author. Department of Oral Biology, Indiana University School of Dentistry, 1121 West Michigan Street, DS 271, Indianapolis, IN 46202, USA. Tel.: +1 317 274 1448; fax: +1 317 278 1411.

E-mail address: [ljwindsor@iu.edu](mailto:ljwindsor@iu.edu) (L.J. Windsor).

Peer review under responsibility of The Center for Food and Biomolecules, National Taiwan University.

Periodontal diseases involve inflammation of the supporting connective tissues with subsequent loss of alveolar bone.<sup>4,5</sup> Overproduction and/or activation of the host derived MMPs eventually lead to the destruction of the periodontal extracellular matrix.<sup>6</sup> MMP-9 can be activated by hypochlorous acid (HOCl) generated by neutrophil myeloperoxidase (MPO) utilizing H<sub>2</sub>O<sub>2</sub> as a substrate.<sup>7</sup> This suggests that ROS production and MMP-mediated periodontal diseases are directly related.<sup>8</sup>

The main polyphenols in Green tea (綠茶 *lù chá*) extract (GTX) are derivatives of catechin (flavanols) that include epigallocatechin gallate (EGCG), epigallocatechin, gallic acid, gallic acid gallate, epicatechin and epicatechin gallate. GTX has anti-oxidant activities. GTX and its catechins have been shown to inhibit ROS production including superoxide (O<sub>2</sub><sup>-</sup>) and nitric oxide (NO) *in vitro* in a dose-dependent manner.<sup>9</sup> Green tea catechins have also been shown to be effective in the prevention and treatment of periodontal disease and dental caries due to their anti-microbial activity.<sup>10,11</sup> Their inhibition of bacterial amylase activity and the growth of oral microorganisms have been documented.<sup>12,13</sup> GTX has also been shown to have anti-inflammatory and anti-proliferative properties.<sup>14,15</sup>

Since the activation and release of MMP-9 from stimulated neutrophils upon infection play a part in inducing periodontal diseases and other chronic inflammatory diseases, any agent that could inhibit the production of ROS and also inhibit the activity or release of MMP-9 from stimulated neutrophils would be a novel “multi-pronged” mechanism for the prevention and treatment of various inflammatory diseases including periodontal disease. In the present study, the ability of GTX and its major catechin, EGCG, to inhibit the activity of MMP-9 and its release from stimulated neutrophils were examined *in vitro*.

## 2. Materials and methods

### 2.1. Reagents

Green tea (*Camellia sinensis*; 綠茶 *lù chá*) was cultivated in the Chonnam province in Korea. The green tea was then stored and processed at the Department of Food and Technology, Chonnam University, Gwangju, Korea. After being air-dried under a fume hood at room temperature, the aerial parts of the plants were crushed using a super mixer (model SM2000, Retch, Germany). The dried materials were extracted in 80% methanol (MeOH) for 24 h and filtered through Whatman No. 2 filter paper. The MeOH extract was then concentrated using a vacuum evaporator (Type N-2N, Eyela, Tokyo, Japan) attached to a cooling aspirator. The concentrated MeOH GTX was then lyophilized (FDU-540, Eyela, Tokyo, Japan). The GTX was then dissolved in 10 mM dimethyl sulphoxide (DMSO) before being diluted in phosphate buffered saline (PBS). The EGCG (95 % purity) was purchased from Calbiochem (La Jolla, CA). EGCG was diluted in PBS prior to conducting the experiments. Human MMP-9 was purified by gelatin sepharose from conditioned media as previously described.<sup>16</sup>

### 2.2. EGCG analysis

High-performance liquid chromatography (HPLC) equipped with a UV-Vis detector was used to determine the EGCG content using maximal absorption peaks at 280 nm. HPLC analysis was performed on Agilent 1100 system (Palo Alto, CA) using an Eclipse XDB-C18 chromatography column (3.5  $\mu$ m, 4.6  $\times$  150 mm) with a 20  $\mu$ L injection volume. A binary mobile phase consisting of solvent systems A and B were used in an isocratic elution with 90:10 A:B. Mobile phase A and B was 2% formic acid (v/v) in ddH<sub>2</sub>O and 100% acetonitrile, respectively. The mobile phase flow rate was 1.0 mL/min and

the run time was 15 min. The retention time for EGCG was 6.07 min. The amount of EGCG was calculated from a standard curve equation.

### 2.3. $\beta$ -Casein cleavage assay

The ability of GTX or EGCG to inhibit the activity of MMP-9 was examined using a  $\beta$ -Casein cleavage assay. Briefly, activated purified samples (3.0  $\mu$ g/ml) of human MMP-9 were incubated with 1.25 mg/ml  $\beta$ -casein (Mr 21,000) at 37 °C.<sup>17</sup> Samples with or without GTX or EGCG were periodically (0–60 min) removed and the reactions stopped by addition of 1, 10-phenanthroline to a final concentration of 10 mM. The samples were analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), stained with Coomassie blue and analyzed by densitometry. Serial dilution of GTX was utilized to determine the dose-response.

### 2.4. Human neutrophils

Buffy coats separated from healthy human donor blood were obtained from the Central Indiana Regional Blood Center (Indianapolis, IN) with Institutional Review Board approval. A double-sucrose gradient, Histopaque-1119 (3 ml) and Histopaque-1077 (3 ml), was used to separate the neutrophils by centrifugation at 20 °C for 35 min. The lower band containing the granulocytes was drawn off. After washing with 10 ml PBS, the cells were centrifuged at 950 rpm for 10 min and the supernatant was discarded. The washing procedure was repeated twice and the cells were resuspended in 10 ml of Roswell Park Memorial Institute (RPMI) 1640 (Sigma Co. St. Louis, MO) media.<sup>18</sup> After counting, the cells were adjusted to  $2.2 \times 10^7$  cells/ml. Trypan Blue staining was utilized to determine the viability of the harvested cells.

### 2.5. Gelatin zymography

To determine if GTX or EGCG could inhibit the release of MMP-9 from the neutrophils, gelatin zymography was utilized. Human neutrophils ( $10^6$ /ml) were stimulated with  $10^{-6}$  M N-formyl-Met-Leu-Phe (FMLP) for 30 min at 37 °C with or without GTX or EGCG. After incubation, the cells were pelleted by centrifugation at 14,000 rpm for 5 min and the collected supernatant was analyzed for MMP-9 release. The supernatant samples were resolved in 10% SDS-PAGE gels co-polymerized with 1 mg/ml gelatin. After electrophoresis, the gels were washed with solution 1 (50 mM Tris, pH 7.5, 3 mM Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, 2.5 % Triton X-100), solution 2 (50 mM Tris, pH 7.5, 3 mM Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, 5 mM CaCl<sub>2</sub>, 1  $\mu$ M ZnCl<sub>2</sub>, 2.5 % Triton X-100) and solution 3 (50 mM Tris, pH 7.5, 3 mM Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, 5 mM CaCl<sub>2</sub>, 1  $\mu$ M ZnCl<sub>2</sub>) for 20 min each. The gels were then incubated in fresh solution 3 overnight at 37 °C. The gels were later stained with Coomassie blue to visualize the lytic bands. The density of the bands was analyzed by NIH 1.62 Image.

### 2.6. Statistical analysis

The significance between the control and test groups was determined by the Student's t-test. The data was expressed as mean  $\pm$  SEM. A value of  $p < 0.05$  was considered statistically significant.

## 3. Results

### 3.1. EGCG analysis

The amount of EGCG was calculated from the standard curve equation ( $r^2 = 0.99917$ ) and the concentration of EGCG in the 1%

Download English Version:

<https://daneshyari.com/en/article/5635403>

Download Persian Version:

<https://daneshyari.com/article/5635403>

[Daneshyari.com](https://daneshyari.com)