

Clinical Paper Dental Implants

Influence of implant location in patients with and without type 2 diabetes mellitus: 2-year follow-up[☆]

T. Abduljabbar¹, F. Javed²,
V. R. Malignaggi³, F. Vohra¹,
S. V. Kellesarian²

¹Department of Prosthetic Dental Sciences, College of Dentistry, King Saud University, Riyadh, Saudi Arabia; ²Department of General Dentistry, Eastman Institute for Oral Health, University of Rochester, New York, USA; ³Department of General Dentistry, Faculty of Dentistry, Universidad Santa Maria, Caracas, Venezuela

T. Abduljabbar, F. Javed, V. R. Malignaggi, F. Vohra, S. V. Kellesarian: Influence of implant location in patients with and without type 2 diabetes mellitus: 2-year follow-up. Int. J. Oral Maxillofac. Surg. 2017; xxx: xxx–xxx. © 2017 International Association of Oral and Maxillofacial Surgeons. Published by Elsevier Ltd. All rights reserved.

Abstract. The aim of the present cross-sectional retrospective 2-year follow-up clinical study was to assess the influence of implant location on clinical and radiographic parameters around dental implants placed in patients with and without type 2 diabetes mellitus (T2DM). Twenty-seven patients with T2DM and 25 non-diabetic controls were included. Implants were classified into three zones according to their location: (1) anterior zone: implant/s replacing anterior teeth, (2) middle zone: implant/s replacing premolars, and (3) posterior zone: implant/s replacing molars. Peri-implant bleeding on probing (BOP), probing depth (PD), and crestal bone loss (CBL) were measured. *P*-values less than 0.05 were considered statistically significant. The mean age of patients with T2DM was 42.5 years and that of non-diabetic controls was 40.6 years. The mean fasting blood glucose levels of patients with and without T2DM were 74.5 mg/dl (66–80 mg/dl) and 82.5 mg/dl (79–88.1 mg/dl), respectively. The mean duration of T2DM was 4.3 years. There was no significant difference in BOP, PD, or CBL around implants placed in any of the zones in the jaws of patients with and without T2DM. There is no influence of implant location on clinical and radiographic parameters around dental implants placed in patients with and without T2DM.

Key words: dental implants; osseointegration; alveolar bone loss; type 2 diabetes mellitus.

Accepted for publication

According to Albrektsson et al., the crestal bone loss (CBL) around dental implants is

a critical outcome variable that determines the overall success of dental implants¹. They reported that a CBL of up to 1.5 mm around the implant followed by a CBL of 0.2 mm annually is considered normal¹.

A variety of local and systemic factors (such as the location of the implant in the jaws and diabetes mellitus, respectively) have been reported to influence CBL around dental implants^{2,3}. In the anterior maxilla, the alveolar process exhibits a

[☆] This work was conducted at Eastman Institute for Oral Health, University of Rochester, NY 14620, USA.

thin labial and thick palatal cortical plate as compared to the posterior maxilla, which has a thicker buccal Plate⁴. Results from a recent cone beam computed tomography study examining the buccal plate thickness of the maxillary and mandibular dentition showed that the buccal bone thickness is significantly greater from the coronal to the apical direction in the mandibular teeth than in the maxillary dentition⁴. Another zone of the alveolar ridge that is associated with vertical bone deficiency is located at the base of the maxillary sinuses. The placement of dental implants in this zone may require adjunct therapeutic protocols such as guided bone regeneration. Furthermore, it is well known that bone quality (type 1 to type 4) also varies among the jaws. According to Truhlar et al., the densest bone exists in the anterior mandible, followed by the posterior mandible, anterior maxilla, and posterior maxilla⁵. These results suggest that the amount of CBL around osseointegrated dental implants will be influenced by the location of the implants in the jaws.

Several studies have reported that chronic hyperglycemia in patients with diabetes mellitus is a significant risk factor for soft tissue inflammation and CBL around osseointegrated implants and teeth^{6–11}. An explanation in this regard is that chronic hyperglycemia has been associated with an increased formation and accumulation of advanced glycation end-products in the systemic and oral tissues, which in turn increase the release of proinflammatory cytokines that enhance CBL around the natural dentition and implants^{12–14}. However, it is pertinent to mention that under optimal glycemic control, dental implants can osseointegrate and remain functionally stable over long durations in diabetic patients in a manner similar to that in non-diabetic controls².

The present cross-sectional retrospective clinical study was based on the hypotheses that (1) peri-implant soft tissue inflammation and CBL are significantly higher in patients with type 2 diabetes mellitus (T2DM) than non-diabetic controls, and (2) peri-implant soft tissue inflammation and CBL around implants placed in patients with T2DM and controls is independent of the location of the implant in the jaws. Therefore, the aim of the present 2-year follow-up study was to assess the influence of implant location on clinical and radiographic parameters around dental implants placed in patients with and without T2DM.

Materials and methods

Ethical guidelines

The study was approved by the Research Ethics Review Committee of the College of Dentistry, King Saud University, Riyadh, Saudi Arabia. An information sheet (describing the purpose of the study) and a consent form were presented to all participants. Consenting individuals were requested to sign the consent form and were given the freedom to withdraw from the study at any stage of the investigation.

Inclusion and exclusion criteria

The inclusion criteria were as follows: (1) individuals who had undergone dental implant therapy; (2) individuals with T2DM; (3) at least 2 years of follow-up; (4) signing of the consent form. Third molars, tobacco and smokeless tobacco users, use of bone grafting techniques, individuals with systemic disorders such as AIDS, cardiovascular disorders, and renal disorders, pregnant/lactating females, and individuals who had consumed antibiotics, non-steroidal anti-inflammatory drugs, and/or corticosteroids within the past 6 months were excluded.

Study design and participants

The present 2-year follow-up clinical study was based on a cross-sectional and retrospective design. In total, 27 patients with T2DM and 25 self-reported non-diabetic controls were included in the present study. Patients with T2DM were requested to present their medical records for verification of the diagnosis of T2DM.

Classification of implants according to their location in the jaws

Depending upon the location of the implant in the maxilla and mandible, the implants were classified into three zones as follows: (1) anterior zone: implant/s replacing anterior teeth (numbers 11–13, 21–23, 31–33, and/or 41–43), (2) middle zone: implant/s replacing premolars (numbers 14, 15, 24, 25, 34, 35, 44, and/or 45), and (3) posterior zone: implant/s replacing molars (numbers 16, 17, 26, 27, 36, 37, 46, and/or 47).

Clinical and radiographic parameters

All participants were requested to visit an oral healthcare clinic in a fasting state for clinical and radiographic evaluation. All clinical and radiographic assessments were performed by a single trained and

calibrated examiner (TA). The kappa value for intra-examiner reliability was 0.91. Peri-implant bleeding on probing (BOP) and probing depth (PD) were measured around all implants placed in patients with and without T2DM. Both BOP and PD were investigated at six sites per implant (mesiobuccal, midbuccal, distobuccal, mesiolingual, midlingual, and distolingual). The long-cone paralleling technique was used to take digital full-mouth radiographs¹⁵. In brief, patients were seated upright with the floor parallel to the Frankfurt horizontal plane. To standardize the angulation between the X-ray beam and the film, a film holder was used (Dentsply Rinn, York, PA, USA). The central X-ray beam was directed perpendicular to the film and long axis of the implant. All radiographs were viewed on a computer screen at 20 × magnification using Corel-Draw 11.0 software (Corel Corp. and Coral Ltd, Ottawa, Canada). For the determination of CBL, the linear distance from the implant platform to the most coronal portion of the alveolar crest (on the mesial and distal surface) was recorded in millimeters¹⁶. CBL was defined as the distance from the widest supracrestal part of the implant to the alveolar crest¹⁷.

Measurement of fasting blood glucose levels

Fasting blood glucose (FBG) levels were measured in patients with and without T2DM at the time of clinical and radiographic examination using a digital glucometer (OneTouch Verio; Johnson & Johnson Co., Milpitas, CA, USA).

Dental prophylaxis

All participants were enrolled in a biannual dental prophylaxis program (every 6 months for 2 years of follow-up), which involved mechanical plaque and calculus removal from all teeth and/or implant surfaces using an ultrasonic scaler (VV Dental, Guangxi, China).

Statistical analysis

The statistical analysis was performed using SPSS version 18 software (SPSS Inc., Chicago, IL, USA). BOP, PD, and CBL were assessed within and between the groups (patients with and without T2DM) using one-way analysis of variance (ANOVA). For multiple comparisons, the Bonferroni post-hoc test was used. A power analysis was performed using computer software nQuery Advisor 5.0 (Statistical Solutions, Saugus, MA,

Download English Version:

<https://daneshyari.com/en/article/5639011>

Download Persian Version:

<https://daneshyari.com/article/5639011>

[Daneshyari.com](https://daneshyari.com)