

# Effect of Sodium Bicarbonate Buccal Infiltration on the Success of Inferior Alveolar Nerve Block in Mandibular First Molars with Symptomatic Irreversible Pulpitis: A Prospective, Randomized Double-blind Study

Masoud Saatchi, DDS, MS,\* Ali Reza Farhad, DDS, MS,\* Naghmeh Shenasa, DDS, MS,<sup>†</sup> and Saeideh Karimi Haghighi, BS, MS<sup>‡</sup>

## Abstract

**Introduction:** The purpose of this prospective, randomized, double-blind study was to evaluate the effect of a buccal infiltration of sodium bicarbonate on the anesthetic success of the inferior alveolar nerve block (IANB) for mandibular first molars in patients with symptomatic irreversible pulpitis. **Methods:** One hundred patients diagnosed with symptomatic irreversible pulpitis of a mandibular first molar were selected. The patients randomly received a buccal infiltration injection of either 0.7 mL 8.4% sodium bicarbonate with 0.3 mL 2% lidocaine containing 1:80,000 epinephrine or 0.7 mL sterile distilled water with 0.3 mL 2% lidocaine containing 1:80,000 epinephrine in a double-blind manner. After 15 minutes, all the patients received conventional IANB injection using 3.6 mL 2% lidocaine with 1:80,000 epinephrine. Access cavity preparation was initiated 15 minutes after the IANB injection. Lip numbness was a requisite for all the patients. Success was determined as no or mild pain on the basis of Heft-Parker visual analog scale recordings upon access cavity preparation or initial instrumentation. Data were analyzed using the *t*, chi-square and Mann-Whitney *U* tests. **Results:** The success rate after the buccal infiltration of sodium bicarbonate was 78%, whereas without the buccal infiltration of sodium bicarbonate it was 44% ( $P < .001$ ). **Conclusions:** A buccal infiltration of 0.7 mL 8.4% sodium bicarbonate increased the success rate of IANBs in mandibular first molars with symptomatic irreversible pulpitis. (*J Endod* 2016; ■:1–4)

## Key Words

Acid-sensing ion channel, inferior alveolar nerve block, irreversible pulpitis, local anesthesia, transient receptor potential vanilloid receptor type 1

The inferior alveolar nerve block (IANB) is the most routine technique used to anesthetize mandibular molars for endodontic treatment (1, 2). However, the success rate is not always adequate to ensure profound pulpal anesthesia, particularly in patients with irreversible pulpitis (3, 4).

The most likely explanation for the decrease in efficacy of local anesthesia in inflamed pulp can be the activation effect of inflammation on the peripheral free terminals of nociceptive neurons and associated central mechanisms (5–9). During inflammation, protons (hydrogen ions) are among the first mediators released by damaged cells, inducing a local pH fall (10, 11). The lowering pH plays a dominant role in the inflammatory activation and sensitization of nociceptive neurons (11–13). This is caused by activation of different ionic channels such as acid-sensing ion channels (ASICs) (11), transient receptor potential channels (8, 11), and tetrodotoxin-resistant sodium channels (14) (Fig. 1). Therefore, it is hypothesized that buccal infiltration of an alkalinizing agent will reduce nociceptor activation by increasing the pH of the inflamed tissue and will result in improved efficacy of local anesthesia in patients with symptomatic irreversible pulpitis.

Sodium bicarbonate is an alkalinizing agent that is commonly used for the treatment of metabolic acidosis. It also has been used for buffering of local anesthetics. Some studies have shown that increasing the pH of a local anesthetic solution by adding sodium bicarbonate reduces the pain of injection (15–18), accelerates the onset of anesthesia (15, 17, 18), and improves the efficacy of anesthesia (18–20). Others have reported that adding sodium bicarbonate to local anesthetic solutions cannot reduce the pain of injection (21–23), accelerate the onset of anesthesia (22, 23), or


## Significance

The administration of a buccal infiltration injection of 0.7 mL 8.4% sodium bicarbonate before an inferior alveolar nerve block injection can be helpful for clinicians to improve the efficacy of the anesthesia in mandibular first molars with symptomatic irreversible pulpitis.

From the \*Dental Research Center, <sup>†</sup>Department of Endodontics, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran; and <sup>‡</sup>Neuroscience Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Address requests for reprints to Dr Naghmeh Shenasa, Department of Endodontics, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran, 81746-73461. E-mail address: [naghmeh.shenasa@gmail.com](mailto:naghmeh.shenasa@gmail.com)  
0099-2399/\$ - see front matter

Copyright © 2016 American Association of Endodontists.  
<http://dx.doi.org/10.1016/j.joen.2016.07.004>



**Figure 1.** The activation effect of acidic extracellular pH on the ion channel receptors. ASICs, acid-sensing ion channels; TRPV1, transient receptor potential vanilloid 1 channel; TTX-Rs, tetrodotoxin-resistant sodium channels.

improve the efficacy of anesthesia (23, 24). Two recent studies have reported that adding sodium bicarbonate to lidocaine cannot improve the efficacy of IANBs in patients with irreversible pulpitis (25, 26).

There are no studies on the use of buccal infiltration injection of an alkalinizing agent for IANBs in teeth with irreversible pulpitis. The purpose of this prospective, randomized, double-blind study was to evaluate the effect of a buccal infiltration injection of 0.7 mL 8.4% sodium bicarbonate on the success rate of IANBs for mandibular first molars in patients with symptomatic irreversible pulpitis.

## Materials and Methods

One hundred healthy adult patients participated in this study. They were emergency patients of the clinic of the endodontic department of Isfahan University of Medical Sciences, Isfahan, Iran. We excluded potential patients who were under 18 years of age, had a history of significant medical conditions, were pregnant, were allergic to local anesthetics or sulfites, were taking any medications that might influence anesthetic assessment, had active sites of pathosis in the area of injection, and were unable to give informed consent. The ethics committee of the university approved the protocol of the study (no. 3941020, ClinicalTrials.gov number, NCT02726737). Written informed consent was obtained from all the patients. To qualify for inclusion in the study, each patient had a vital mandibular first molar with active moderate to severe pain and had a prolonged response to cold testing with cold spray (Endo-Frost; Coltène-Whaledent, Langenau, Germany). Patients with no response to cold testing, no vital coronal pulp tissue on access, or a periapical lesion (other than widening of the periodontal ligament space) were not included in the study. Therefore, each patient had a vital mandibular first molar with a clinical diagnosis of symptomatic irreversible pulpitis.

Each patient rated his or her initial pain on a Heft-Parker visual analog scale (HP-VAS) (27). This scale is a 170-mm marked line that is divided into 4 categories with various terms describing the level of pain. No pain, mild pain, moderate pain, and severe pain were indicated by 0 mm, 1- to 54-mm, 55- to 113- mm, and 114- to 170-mm divisions, respectively. Patients exhibiting moderate to severe initial pain were included in the study.

For the sodium bicarbonate group, 0.7 mL 8.4% sodium bicarbonate (Caspian Tamin Pharmaceutical Co, Rasht, Iran) was drawn with a sterile microliter syringe (Hamilton, Bonaduz, Switzerland) fitted

with a disposable 27-G needle. Then, 0.3 mL 2% lidocaine with 1:80,000 epinephrine (Lignospan; Septodont, Saint Maur des Fosses, France) was drawn with the same syringe. Therefore, the sodium bicarbonate solution contained 0.7 mL 8.4% sodium bicarbonate and 0.3 mL 2% lidocaine. The syringe was inverted 5 times to mix the solution, and no precipitation was observed. For the non-sodium bicarbonate group, 0.7 mL sterile distilled water was drawn with the Hamilton syringe. Then, 0.3 mL 2% lidocaine with 1:80,000 epinephrine was added in the same way. Therefore, the non-sodium bicarbonate solution contained 0.7 mL sterile distilled water and 0.3 mL 2% lidocaine. A trained dental assistant prepared the solutions just before the injections and coded them in a random manner. One operator administered buccal infiltration injections of either sodium bicarbonate or non-sodium bicarbonate solution for each patient. The operator and patients were both blinded to the contents of the solution.

Before each buccal infiltration, the mucosa was dried, and 20% benzocaine gel (Ultradent Products Inc, South Jordan, UT) was applied to the injection site for 60 seconds. The intended target site was centered over the buccal root apices of the mandibular first molar to be treated. The 27-G needle was placed into the alveolar mucosa and advanced until the needle was estimated to be at or just superior to the apices of the tooth. The solution was deposited over a period of 1 minute.

After 15 minutes, the same operator administered 2 standard IANBs (1.8-mL cartridges) of 2% lidocaine with 1:80,000 epinephrine (Lignospan; Septodont, Saint Maur des Fosses, France) for each patient. All the injections were performed using a standard aspirating dental injection syringe and a 27-G, 31-mm needle (CK ject; CK Dental, Kor-Kyungji-do, Korea). Lip numbness was considered as a criterion for IANB achievement; the patient was questioned for lip numbness 15 minutes after the injection. If lip numbness was not achieved, the IANB was indicated as missed, and the patient was excluded from the study. No patient was excluded from the study as a result of a lack of lip numbness. Fifteen minutes after the injection, the teeth were isolated with a rubber dam, and access cavities were prepared.

The same operator instructed the patients to rate any pain felt during access cavity preparation or the initial file placement. If the patient felt pain, the treatment was immediately ceased, and the patient rated the pain level by using the HP-VAS. The success of the IANB was defined as no pain or mild pain (HP-VAS score  $\leq 54$ ).

## Statistical Analysis

Data on age, sex, initial pain, and the success of IANB ratings were statistically analyzed using SPSS software version 20 (IBM Corporation, Armonk, NY). Comparisons between the sodium bicarbonate and non-sodium bicarbonate groups for the success of the IANB and sex differences were analyzed using the chi-square test, age was analyzed using the *t* test, and initial pain was analyzed using the Mann-Whitney *U* test. With a 2-sided alpha risk of 0.05, a sample size of 50 subjects per group was required to detect a difference of  $\pm 30$  percentage points in anesthetic success with a power of more than 0.80. Statistical significance was defined as  $P < .05$ .

## Results

One hundred adult patients, 27 men and 73 women, with an age range of 18–53 years and a mean of  $29 \pm 9$  years, participated in the study. Baseline variables for the sodium bicarbonate and non-sodium bicarbonate groups are presented in Table 1. There were no significant differences in age, sex, or initial pain between the 2 groups ( $P > .05$ ). The IANB success rate was 78% for the sodium bicarbonate group and 44% for the non-sodium bicarbonate group. There was a statistically significant difference in success rates between the 2 groups ( $P < .001$ ).

Download English Version:

<https://daneshyari.com/en/article/5641062>

Download Persian Version:

<https://daneshyari.com/article/5641062>

[Daneshyari.com](https://daneshyari.com)