

# Postoperative Delirium Following Orthognathic Surgery in a Young Patient



*Fernanda Herrera da Costa, DMD, \* Paulo Adilson Herrera, MD, MSc, †  
Cecília Luiz Pereira-Stabile, DMD, MSc, PhD, ‡ and  
Glaykon Alex Vitti Stabile, DMD, MSc, PhD §*

Delirium is an organic mental syndrome with acute onset characterized by diffuse brain dysfunction and neural activity disorganization. It is usually related to cognition and perception changes, decreased level of consciousness, and disorganization of thoughts that are unrelated to previous dementia. Occurrence is more frequent in patients with previous degenerative disease and elderly patients, especially those older than 85 years. Although the pathophysiology is not totally known, studies have shown that, among the main factors that lead to delirium, the drugs used for general anesthesia induction are the most relevant (hypnotics, anticholinergic drugs, and H<sub>2</sub> receptor blockers), especially those used in long surgical procedures. This report describes the case of a 24-year-old woman with a noncontributory medical and psychological history. She underwent bimaxillary orthognathic surgery with a total general anesthesia time of 7 hours. Postoperatively, she developed agitation, confusion, and delirium. After a psychiatry consult and discussion with the anesthesia team, the diagnosis of psychotic break owing to late postoperative delirium was established.

© 2016 Published by Elsevier Inc on behalf of the American Association of Oral and Maxillofacial Surgeons

*J Oral Maxillofac Surg* 75:284.e1-284.e4, 2017

Orthognathic surgery is a complex and specialized procedure that might require long periods of general anesthesia. Postoperative delirium (POD) is a common complication involving postoperative disturbance of the mental state, especially in patients older than 85 years, with an incidence of 15 to 53% in the elderly population.<sup>1</sup> However, the number of occurrences increases as the procedure lengthens, such as heart and orthopedic procedures,<sup>2</sup> with an even higher incidence when associated with pre-existing mental disorders.<sup>3</sup> School-age children appear to be at risk, with an incidence of 12% when undergoing eye surgery. Its occurrence in adolescents and young adults is rare.<sup>4</sup> In addition to age, risk factors possibly associated

with POD are Parkinson disease, malnutrition, head trauma, prolonged anesthesia, infection, and metabolic disorders.<sup>1,5</sup>

POD is an acute-onset mental syndrome characterized by global cognitive dysfunction: a lower level of consciousness, abnormal attention, increase or decrease in psychomotor activity, and dysfunction of the sleep-and-wake cycle.<sup>6</sup> The etiology is multifactorial, and the most accepted theory is that occurrence is related to changes in neurotransmission caused by drugs, neurologic conditions, underlying diseases, environmental factors, trauma, or longer exposure to general anesthesia.<sup>2,7</sup> The drugs most commonly involved in the development of delirium are

\*Surgeon, Department of Oral and Maxillofacial; MSc Student, Department of Oral Pathology, State University of São Paulo, São Paulo, SP, Brazil.

†Private Anesthesiologist, Hospital Evangélico de Londrina, Londrina, PR, Brazil.

‡Associate Professor, Department of Oral Medicine and Pediatric Dentistry, State University of Londrina, Londrina, PR, Brazil.

§Adjunct Professor, Department of Oral Medicine and Pediatric Dentistry; Coordinator, Residency in Oral and Maxillofacial Surgery, State University of Londrina, Londrina, PR, Brazil.

Address correspondence and reprint requests to Dr Costa: Pernambuco Street, 540, Londrina, PR CEP 86020-120, Brazil; e-mail: [fer\\_odontouel@hotmail.com](mailto:fer_odontouel@hotmail.com)

Received October 11 2015

Accepted August 14 2016

© 2016 Published by Elsevier Inc on behalf of the American Association of Oral and Maxillofacial Surgeons

0278-2391/16/30758-3

<http://dx.doi.org/10.1016/j.joms.2016.08.024>

narcotics, hypnotics, H<sub>2</sub> receptor blockers, drugs for the treatment of Parkinson disease, anticholinergic agents, and corticosteroids. The potential for drugs to induce delirium depends on underlying risk factors and individual susceptibility.<sup>1,8</sup> Thus far, the occurrence of POD in association with orthognathic surgery has not been reported in the literature.

Through the report of a clinical case, the aim of this article was to describe and characterize POD as a possible complication of orthognathic surgery.

## Report of Case

This study was approved by the institutional review board of the Hospital Evangélico de Londrina (Londrina, Brazil) and the patient signed an informed consent agreement.

A 24-year-old woman was referred by her orthodontist to the oral and maxillofacial surgery team for an orthognathic evaluation. She denied having any health issues except for a diagnosis of von Willebrand disease. The patient was professionally active in the preoperative period and did not present with any mental disorders or systemic comorbidities. Two hours before surgery, as recommended by her hematologist, she received a transfusion of coagulation factor VIII. As a preanesthetic drug, midazolam 15 mg was taken orally. Upon arrival in the operating room, cefazolin 1 g and dexamethasone 10 mg were administered intravenously. Anesthesia was induced by intravenous injection of propofol, remifentanyl, and rocuronium, maintained by isoflurane inhalation and continuous infusion of remifentanyl, and monitored by a mass spectrometer. During the procedure, the patient was continuously monitored by electrocardiography, radial artery puncture, capnography, and a bispectral index (BIS) monitor. Mean arterial pressure was maintained at 65 to 80 mmHg with the aid of a  $\beta$ -blocker (metoprolol) and the BIS was kept at 40 to 60. During the procedure, the patient's electroencephalogram did not show any suppression periods. There was no need for blood transfusion during surgery or the postoperative period.

After anesthetic induction, the patient underwent orthognathic surgery for correction of a Class II dento-facial deformity through maxillary impaction, mandibular advancement, and an advancement genioplasty. Surgery was uneventful. The patient remained under general anesthesia for 7 hours.

After extubation and postanesthetic recovery, the patient was transferred to the floor, where she was prescribed an intravenous antibiotic (cephalothin), a corticosteroid (hydrocortisone), and 2 nonsteroidal anti-inflammatory and analgesic medications (ketorolac and dipyrone).

During the first 24 hours after surgery, the patient was calm, oriented in time and space, communicative,

and responsive to verbal commands. She did not exhibit postoperative hemorrhage, was able to take liquids orally, had spontaneous diuresis, moderate swelling, and adequate pain control, and was discharged home the day after surgery. Upon discharge, she was prescribed an oral antibiotic and nonsteroidal analgesic. No further steroids were prescribed.

Within the first 48 hours, the patient's parents contacted the surgical team reporting unusual behavior. She alternated periods of agitation and aggressiveness and was unable to sleep. As time passed, the symptoms became more severe and she became incoherent, presenting delusions of control and thought insertion. She also reported an inability to breathe during sleep and constant thoughts about dying. At the same time, her parents observed an increase in aggressiveness. As a result, the patient was not sleeping adequately and eating precariously.

She was referred for psychiatric emergency services. After rigorous clinical evaluation concurrently with the anesthesiology team, the patient was diagnosed with a psychotic break caused by POD, described in the *International Classification of Diseases, Tenth Revision* as a psychotic disorder from the use of multiple drugs and other psychoactive substances (code F19.5).<sup>9</sup> Treatment consisted of:

- Instructing the caregivers not to leave the patient alone while the symptoms were present
- Prescription of antidepressant drugs (escitalopram and mirtazapine) and an anxiolytic (diazepam), with instructions to increase the dose in case of worsening symptoms

Within the first 24 hours after starting the prescribed medication, the patient showed considerable improvement of mood and signs, and symptoms of paranoid delirium became less severe.

After 3 days of using psychiatric medication, the patient was completely asymptomatic and the drugs were gradually discontinued. At 10 months after surgery, she was totally asymptomatic and resumed all her professional and social activities uneventfully.

## Discussion

The purpose of this report was to characterize POD as a possible complication of orthognathic surgery. According to Lipowski,<sup>6</sup> POD can be classified by its time of occurrence as emergence delirium, when it occurs within the first 24 hours after anesthesia, and late delirium, when it occurs after a lucid interval of at least 1 day.<sup>10</sup> Other investigators have classified POD, based on its clinical presentation, as agitation or hyperactivity, characterized by an increased state of alertness to external stimuli, or silent or

Download English Version:

<https://daneshyari.com/en/article/5642038>

Download Persian Version:

<https://daneshyari.com/article/5642038>

[Daneshyari.com](https://daneshyari.com)