Accepted Manuscript

The increased risk of stroke in early insomnia following traumatic brain injury: a population-based cohort study

Kam-Hou Ao, Chung-Han Ho, Che-Chuan Wang, Jhi-Joung Wang, Chung-Ching Chio, Jinn-Rung Kuo

PII: \$1389-9457(17)30089-8

DOI: 10.1016/j.sleep.2017.02.010

Reference: SLEEP 3316

To appear in: Sleep Medicine

Received Date: 5 September 2016

Revised Date: 6 January 2017 Accepted Date: 9 February 2017

Please cite this article as: Ao K-H, Ho C-H, Wang C-C, Wang J-J, Chio C-C, Kuo J-R, The increased risk of stroke in early insomnia following traumatic brain injury: a population-based cohort study, *Sleep Medicine* (2017), doi: 10.1016/j.sleep.2017.02.010.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1

The increased risk of stroke in early insomnia following traumatic brain injury: a population-based cohort study

Kam-Hou Ao ^{a,*}, Chung-Han Ho ^{b,c},*, Che-Chuan Wang ^{a,d}, Jhi-Joung Wang ^b, Chung-Ching Chio ^a, Jinn-Rung Kuo ^{a,b,e,†}

Departments of ^a Neurosurgery and ^b Medical Research, Chi-Mei Medical Center, Tainan, Taiwan

^c Department of Pharmacy, Chia Nan University of Pharmacy and Science, Tainan, Taiwan

Departments of ^e Biotechnology, and ^d Child Care, Southern Taiwan University of Science and Technology, Tainan, Taiwan

* These authors contributed equally to this work.

[†] Corresponding author. Department of Neurosurgery, Traumatic Brain Center, Chi-Mei Medical Center, #901, Jong-Hwa Road, Yung-Kung district, Tainan City 710, Taiwan. Tel.: +886 6 281 2811 ext. 9; fax: +886 6 282 8928.

E-mail address: kuojinnrung@gmail.com (J.-R. Kuo).

ABSTRACT

Objective: Insomnia, a common symptom after traumatic brain injury (TBI), may be a pre-symptom for developing stroke. This study aims to investigate whether insomnia is a potential risk factor for stroke after TBI, especially early insomnia.

Download English Version:

https://daneshyari.com/en/article/5643878

Download Persian Version:

https://daneshyari.com/article/5643878

<u>Daneshyari.com</u>