



# Isolated subcutaneous sarcoid-like granulomatous inflammation occurring at injection sites: 3 patients treated successfully with minocycline

James Abbott, BS,<sup>a</sup> Laura A. Taylor, MD,<sup>a</sup> Karolyn A. Wanat, MD,<sup>b</sup> and Misha Rosenbach, MD<sup>c</sup>  
*Philadelphia, Pennsylvania and Iowa City, Iowa*

**Key words:** desensitizing agents; granulomatous inflammation; leuprolin acetate; minocycline; sarcoidal granulomas; sarcoidosis; subcutaneous granulomas; subcutaneous sarcoidosis.

## INTRODUCTION

Subcutaneous sarcoidosis is a rare manifestation of systemic sarcoidosis in which firm asymptomatic-to-tender subcutaneous nodules develop secondary to sarcoidal granulomas involving the panniculus.<sup>1</sup> Sarcoidosis implies systemic disease, but subcutaneous lesions may present with absent or mild systemic involvement. Here we report 3 cases of isolated subcutaneous nodules with sarcoidal granulomas after subcutaneous injections that improved after treatment with minocycline.

## REPORT OF A CASE SERIES

In all 3 patients, sarcoidal granulomas developed at the site of previous iatrogenic injections (allergy shots and leuprolin acetate) in the absence of additional lesions or systemic manifestations. All 3 patients had lesions on their bilateral posterior arms and had a complete response to minocycline. Relevant clinical information, diagnostic workup, and micrographs are presented in [Table I](#) and [Fig 1](#), respectively.

## DISCUSSION

Subcutaneous nodules are rare cutaneous manifestations of sarcoidosis occurring in up to 6% of cases, frequently manifesting on the forearms as

indurated linear bands, although lesions can occur at any site.<sup>1-3</sup> In our patients, lesions developed on the bilateral posterior arms, which correspond to prior inoculation sites and differ from the common locations of subcutaneous sarcoidosis.

Previous reports show that subcutaneous sarcoidosis may be the initial clinical presentation of mild systemic involvement.<sup>2,3</sup> However, because of the isolated nature of many cases, it is debated if these subcutaneous granulomatous lesions truly represent a sarcoidal variant. Nevertheless, these lesions require workup for systemic disease and treatment for physical discomfort and aesthetics.

Although the exact etiology of sarcoidosis remains elusive, the development of granulomas has been attributed to a hypersensitivity reaction with a nonspecific response to extrinsic or intrinsic antigens in a genetically susceptible individual. Exposure to these antigens produces local granulomas and is postulated to initiate systemic disease.<sup>4</sup>

The treatment for subcutaneous sarcoidosis revolves around systemic corticosteroids and immunosuppressive agents.<sup>2</sup> In our patients, a therapeutic response was seen with minocycline as a first-line therapy. All patients had a complete clinical resolution; the 2 patients who had preradiographic and postradiographic studies showed

From the Departments of Pathology<sup>a</sup> and Dermatology,<sup>c</sup> The University of Pennsylvania School of Medicine and the Department of Dermatology and Pathology, University of Iowa College, Hospital and Clinics.<sup>b</sup>

Funding sources: None.

Conflicts of interest: None declared.

Correspondence to: Misha Rosenbach, MD, Department of Dermatology, Hospital of the University of Pennsylvania, 3600 Spruce St, 2 Maloney Building, Philadelphia, PA 19104. E-mail: [Misha.Rosenbach@uphs.upenn.edu](mailto:Misha.Rosenbach@uphs.upenn.edu).

JAAD Case Reports 2017;3:74-7.

2352-5126

© 2017 by the American Academy of Dermatology, Inc. Published by Elsevier, Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<http://dx.doi.org/10.1016/j.jidcr.2016.12.004>

**Table I.** Clinical features and treatments of cases

	Case 1	Case 2	Case 3
Age/sex	30s/male	50s/female	50s/female
Clinical manifestation	6 tender subcutaneous nodules with papules and plaques	Non-tender subcutaneous nodules; right arm involving 1-1.5% BSA	Tender subcutaneous nodules
Location	Bilateral posterior arms	Bilateral posterior arms	Bilateral posterior arms
Subcutaneous injection	Allergy shots (>20 years ago)	Allergy shots (20-30 years ago)	Leuprolin acetate; 1-2 weeks prior to development of lesions
PMH	Noncontributory	Alpha-1-anti-trypsin (MS genotype)	Uterine fibroids, hypothyroid
ACE (U/L)	97 U/L	39 U/L	14 U/L
Vitamin D 1,25-dihydroxy (pg/mL)	79	N/A	N/A
Vitamin D 25-hydroxy (ng/mL)	25	35	N/A
Calcium (mg/dL)	9.6	9.2	9.7
Chest radiograph	Negative	Negative	Negative
Ophthalmology	Normal	N/A	N/A
Magnetic resonance imaging	None	Interdigitating lesion in the subcutaneous fat that measured 5.0 × 4.9 × 3.0 cm and abuts the superficial muscle fascia	Lobulated soft tissue mass of the left deltoid measuring 3.1 × 1.9 × 4.9 cm, extending beyond the fascia with suspicion for malignancy
Pathology	Nonnecrotizing granulomatous inflammation involving the subcutaneous tissue	Granulomatous panniculitis with sarcoidal-type granulomas	Noncaseating granulomatous inflammation involving the subcutaneous adipose tissue
Treatment	Minocycline	Minocycline	Minocycline
Duration of treatment	2 months; discontinued due to transient muscle pain	2 months; no adverse events	5 months; no adverse events
Treatment response	Complete resolution	Complete resolution	Complete resolution

All cases had additional therapeutic workup, but only the pertinent diagnostics to rule out systemic sarcoidosis were included. ACE, Angiotensin converting enzyme; N/A, not applicable; PMH, past medical history.

Download English Version:

<https://daneshyari.com/en/article/5646258>

Download Persian Version:

<https://daneshyari.com/article/5646258>

[Daneshyari.com](https://daneshyari.com)