


Accepted Manuscript

Title: Modyfied wedge osteotomy for osteoarthritis of elbow secondary to osteochondritis dissecans in adolescent with multiple epiphyseal dysplasia. A case report.

Authors: Takashi Yoshida, Wook-Cheol Kim, Yoshinobu Oka, Masashi Nakase, Atsushi Nishida, Hiroaki Wada, Yuji Arai, Toshikazu Kubo



PII: S0972-978X(17)30188-5
DOI: <http://dx.doi.org/doi:10.1016/j.jor.2017.08.011>
Reference: JOR 405

To appear in:

Received date: 29-5-2017
Accepted date: 6-8-2017

Please cite this article as: Takashi Yoshida, Wook-Cheol Kim, Yoshinobu Oka, Masashi Nakase, Atsushi Nishida, Hiroaki Wada, Yuji Arai, Toshikazu Kubo, Modyfied wedge osteotomy for osteoarthritis of elbow secondary to osteochondritis dissecans in adolescent with multiple epiphyseal dysplasia. A case report. (2010), <http://dx.doi.org/10.1016/j.jor.2017.08.011>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Modified wedge osteotomy for osteoarthritis of elbow secondary to osteochondritis dissecans in adolescent with multiple epiphyseal dysplasia. A case report.

Takashi Yoshida¹, Wook-Cheol Kim¹, Yoshinobu Oka¹, Masashi Nakase¹, Atsushi Nishida¹, Hiroaki Wada¹, Yuji Arai², Toshikazu Kubo¹

Academic affiliations:

¹Department of Orthopaedics, Graduate School of Medical Science, Kyoto Prefectural University of Medicine, Kyoto, Japan

²Department of Sports Para-Sports Medicine, Graduate School of Medical Science, Kyoto Prefectural University of Medicine, Kyoto, Japan

Corresponding author: Takashi Yoshida, M.D., Ph.D.

e-mail: mohi72@koto.kpu-m.ac.jp

Address: 465 Kajii-cho Kawaramachi-Hirokoji, Kamigyo-ku, Kyoto 602-0841, Japan

tel.: +81-75-251-5549, fax.: +81-75-251-5841

Abstract

Multiple epiphyseal dysplasia (MED) may be complicated by osteochondritis dissecans (OCD), most commonly reported in the knee, but involvement of the elbow is very rare. Optimal

Download English Version:

<https://daneshyari.com/en/article/5654106>

Download Persian Version:

<https://daneshyari.com/article/5654106>

[Daneshyari.com](https://daneshyari.com)