
## **Accepted Manuscript**

Development and Validation of a Risk Scoring System for Severe Acute Lower Gastrointestinal Bleeding

Tomonori Aoki, Naoyoshi Nagata, Takuro Shimbo, Ryota Niikura, Toshiyuki Sakurai, Shiori Moriyasu, Hidetaka Okubo, Katsunori Sekine, Kazuhiro Watanabe, Chizu Yokoi, Mikio Yanase, Junichi Akiyama, Masashi Mizokami, Naomi Uemura



PII: S1542-3565(16)30308-1 DOI: 10.1016/j.cgh.2016.05.042

Reference: YJCGH 54795

To appear in: Clinical Gastroenterology and Hepatology

Accepted Date: 31 May 2016

Please cite this article as: Aoki T, Nagata N, Shimbo T, Niikura R, Sakurai T, Moriyasu S, Okubo H, Sekine K, Watanabe K, Yokoi C, Yanase M, Akiyama J, Mizokami M, Uemura N, Development and Validation of a Risk Scoring System for Severe Acute Lower Gastrointestinal Bleeding, *Clinical Gastroenterology and Hepatology* (2016), doi: 10.1016/j.cgh.2016.05.042.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

All studies published in Clinical Gastroenterology and Hepatology are embargoed until 3PM ET of the day they are published as corrected proofs on-line. Studies cannot be publicized as accepted manuscripts or uncorrected proofs.

## ACCEPTED MANUSCRIPT

Development and Validation of a Risk Scoring System for Severe Acute Lower Gastrointestinal Bleeding

**Short title:** Predictors of severe LGIB

Authors: Tomonori Aoki<sup>1</sup>, Naoyoshi Nagata<sup>1</sup>, Takuro Shimbo<sup>2</sup>, Ryota Niikura<sup>3</sup>, Toshiyuki Sakurai<sup>1</sup>, Shiori Moriyasu<sup>1</sup>, Hidetaka Okubo<sup>1</sup>, Katsunori Sekine<sup>1</sup>, Kazuhiro Watanabe<sup>1</sup>, Chizu Yokoi<sup>1</sup>, Mikio Yanase<sup>1</sup>, Junichi Akiyama<sup>1</sup>, Masashi Mizokami<sup>4</sup>, and Naomi Uemura<sup>5</sup>

<sup>1</sup>Department of Gastroenterology and Hepatology, National Center for Global Health and Medicine, 1-21-1 Toyama, Shinjuku, Tokyo 162-8655, Japan

<sup>2</sup>Ohta Nishinouchi Hospital, 2-5-20 Nishinouchi, Koriyama, Fukushima 963-8022, Japan

<sup>3</sup>Department of Gastroenterology, Graduate School of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-8655, Japan

<sup>4</sup>The Research Center for Hepatitis and Immunology, National Center for Global Health and Medicine, Kohnodai Hospital, 1-7-1 Kohnodai, Ichikawa City, Chiba 272-8516, Japan

<sup>5</sup>Department of Gastroenterology and Hepatology, National Center for Global Health and Medicine, Kohnodai Hospital, 1-7-1 Kohnodai, Ichikawa City, Chiba 272-8516, Japan

Grant Support: This study was partly supported by a Grant-in-Aid for research and development from the National Center for Global Health and Medicine (26A-201). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abbreviations: AUC, area under the receiver operating characteristic curve; CI, confidence interval; LGIB, lower gastrointestinal bleeding; LOS, length of stay; MDCT, multidetector computed tomography; NCGM,

## Download English Version:

## https://daneshyari.com/en/article/5657470

Download Persian Version:

https://daneshyari.com/article/5657470

<u>Daneshyari.com</u>