Accepted Manuscript

Should we all be using the Doppler endoscopic probe in non-variceal upper gastrointestinal bleeding?

Jennifer Nayor, MD, John R. Saltzman, MD

A Controlled Trial of Gluten-Free Diet in Museum State Installed State Install

PII: S0016-5085(17)30330-X DOI: 10.1053/j.gastro.2017.03.021

Reference: YGAST 61053

To appear in: Gastroenterology

Please cite this article as: Nayor J, Saltzman JR, Should we all be using the Doppler endoscopic probe in non-variceal upper gastrointestinal bleeding?, *Gastroenterology* (2017), doi: 10.1053/j.gastro.2017.03.021.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Should we all be using the Doppler endoscopic probe in non-variceal upper gastrointestinal bleeding?

Jennifer Nayor, MD^{1,2}, John R. Saltzman, MD^{1,2}

- 1. Division of Gastroenterology, Hepatology and Endoscopy, Brigham and Women's Hospital, Boston, MA
- 2. Harvard Medical School, Boston, MA

The authors have no conflicts of interest.

Corresponding Author:

John R. Saltzman, MD
Director of Endoscopy
Brigham and Women's Hospital
Division of Gastroenterology, Hepatology and Endoscopy
Professor of Medicine
Harvard Medical School
75 Francis Street
Boston, MA 02115

Tel: 617-525-8763 Fax: 617-264-5264

E-mail: jsaltzman@bwh.harvard.edu

Download English Version:

https://daneshyari.com/en/article/5658531

Download Persian Version:

https://daneshyari.com/article/5658531

Daneshyari.com