Accepted Manuscript

Acyl/free carnitine ratio is a risk factor for hepatic steatosis after pancreateduodenectomy and total pancreatectomy

Masafumi Nakamura, Kohei Nakata, Hideo Matsumoto, Takao Ohtsuka, Koji Yoshida, Shoji Tokunaga, Keisuke Hino

PII: \$1424-3903(16)31247-9

DOI: 10.1016/j.pan.2016.11.007

Reference: PAN 652

To appear in: Pancreatology

Received Date: 25 April 2016


Revised Date: 28 November 2016 Accepted Date: 29 November 2016

Please cite this article as: Nakamura M, Nakata K, Matsumoto H, Ohtsuka T, Yoshida K, Tokunaga S, Hino K, Acyl/free carnitine ratio is a risk factor for hepatic steatosis after pancreatoduodenectomy and total pancreatectomy, *Pancreatology* (2016), doi: 10.1016/j.pan.2016.11.007.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT



Download English Version:

https://daneshyari.com/en/article/5661535

Download Persian Version:

https://daneshyari.com/article/5661535

<u>Daneshyari.com</u>