

Video case report

A step-by-step surgical technique video with two reported cases of common channel lengthening in patients with previous stomach intestinal pylorus sparing surgery to treat chronic diarrhea

Amit Surve, M.D., Hinali Zaveri, M.D., Daniel Cottam, M.D.*

Bariatric Medicine Institute, Salt Lake City, UT

Received September 13, 2016; accepted October 15, 2016

Keywords: Common channel lengthening; Stomach intestinal pylorus sparing surgery; Chronic diarrhea; SIPS; Technique; Loop duodenal switch

Surgical procedures are currently the only effective long-term way to treat morbid obesity. However, these procedures all carry risks, with each procedure having its own unique risks.

In 2007, Dr. Sanchez in Spain first performed a modification of the traditional duodenal switch using a single anastomosis instead of Roux-en-Y reconstruction with the sleeve done over a 56 French bougie [1]. We have modified this further by creating a smaller sleeve (40 French bougie sizing instead of 56) and less malabsorption (300-cm common channel instead of 250 cm) [2]. We have called this modification “stomach intestinal pylorus sparing” (SIPS) surgery to help differentiate it from the common complication of the traditional Roux-en-Y duodenal switch. The preservation of more intestine, along with the ileocecal valve, reduces the risk for malnutrition and diarrhea often associated with the duodenal switch [3,4]. The lack of a distal small bowel enteroenterostomy should eliminate or decrease the chances of internal hernias. Since 2013, we have performed over 300 SIPS surgeries. However, we have encountered 2 patients that developed chronic diarrhea without malnutrition that resulted in the need for common channel lengthening. Chronic diarrhea was defined as loose stools that last for at least 4 weeks. This usually means 4 or more loose stools per day.

We present a step-by-step surgical technique of common channel lengthening (CCL) to treat chronic diarrhea after SIPS surgery. This is the first report in the literature that demonstrates the CCL technique after a SIPS surgery and also reports the first 2 cases.

Patient 1

A 41-year-old woman with a body mass index (BMI) of 42.9 kg/m² had planned SIPS in 2014. Her current BMI is 26.5 kg/m². Six months after surgery, she began to experience smelly flatulence, bloating, and severe diarrhea (6–8 loose stools/d). Her nutritional status (vitamins A, D, E, K, B1, B12, and copper and zinc) was normal. Her albumin and prealbumin were also normal. Dietary intervention and medical interventions of Lomotil and probiotics did not help.

Patient 2

The second patient was a 63-year-old man with a history of inflammatory bowel syndrome who underwent SIPS surgery a year prior for morbid obesity with insulin dependent diabetes and a BMI of 38.6 kg/m². The patient had remarkable weight loss after the SIPS surgery (current BMI, 22.5 kg/m²) with resolution of pre-existing diabetes and gastroesophageal reflux disease. However, 5 months after surgery the patient began experiencing postprandial diarrhea (6–8 loose stools/d) that was unresponsive to both dietary manipulation and medical interventions of

*Correspondence: Daniel Cottam, M.D., Bariatric Medicine Institute, 1046 East 100 South, Salt Lake City, UT 84102.
E-mail: drdanielcottam@yahoo.com

probiotics and Lomotil. His protein and nutritional statuses were also normal (albumin, prealbumin, vitamins A, D, E, K, B1, B12, and copper and zinc).

Methods

Data from the patients were drawn from a prospectively kept database. From September 2013 to August 2016, our center performed over 300 laparoscopic SIPS surgeries. Additionally, there were over 50 revisions in which gastric bypass or laparoscopic bands were converted to SIPS cases [5,6]. Two of our patients experienced chronic diarrhea following a SIPS surgery. We elected to perform laparoscopic CCL to increase the absorptive capacity.

The SIPS technique

Our SIPS technique has been described in detail in previous work [2,7].

Management

The CCL technique

Laparoscopic CCL was performed using a standard 4-port technique. Initial access is obtained by means of a Veress needle technique at the left anterior subcostal port site because this site is generally safe from visceral injury. A pneumoperitoneum was established to a pressure of 15 mm Hg using CO₂ gas. A 10-mm 30° laparoscopic camera was used. A liver retractor was placed through the posterior right subcostal port to elevate the left lobe of the liver.

The surgeon (standing on the right side) operates through the right upper abdominal port and right anterior subcostal port. The assistant surgeon (standing on the left side) operates through the 2 left subcostal ports.

Surgical technique

After positioning and port placement, the abdomen was inspected and adhesions were lysed with blunt and sharp dissection as needed.

Construction of enteroenterostomy

In both cases, the first step of this technique was to locate the duodenoileostomy (DI). The efferent limb (common channel) was traced 300 cm from the DI to the ileocecal valve (Fig. 1 A, step 1). Also, 400 cm from the DI to the ligament of Treitz (afferent limb/biliopancreatic limb) was counted out (Fig. 1 A, step 2). A side-to-side enteroenterostomy was performed 15 cm below the DI using a 2 gastrointestinal anastomosis stapler technique (Fig. 1 A, step 3). This step is performed to ensure that, if there is constriction and a division of the small bowel, an obstruction does not form.

Transection of DI. A complete transection was performed at the DI using a gastrointestinal anastomosis stapler (Fig. 1 A, step 4).

Creation of new loop (duodenojejunoscopy)

We elected to lengthen the common channel by 200 cm. To do this, we traced 200 cm proximal to the new enteroenterostomy on the afferent limb (biliopancreatic limb) (Fig. 1 A, step 5).

The new loop limb was brought up and the antimesenteric border of the jejunum was attached to the end of the proximal duodenum staple line using a 2.0 Polysorb (Covidien Ltd., Dublin, Ireland). The loop was set up so the efferent limb (approximate 500 cm) was descending on the patient's right and the afferent limb was ascending from the left. Approximately 2-cm duodenotomy and enterotomy were made, and the enterotomy was closed with a running posterior layer and running anterior layer using 3.0 Polysorb (Covidien).

Antiobstruction stitch and antivolvulus stitch

Another 2 interrupted sutures were placed—1 from the afferent limb to the antrum and the other from the afferent limb to the omentum to prevent chronic nausea and volvulus, respectively (Fig. 1 A, steps 6 and 7) [5]. The anastomosis was tested intraoperatively for leaks (using an endoscope). Skin incisions were closed.

Postoperative care. The patients were allowed sips of water on the evening of surgery and full liquids on postoperative day 1. Before discharge, both patients underwent an upper gastrointestinal series to rule out leak. Patients were discharged when oral intake was adequate and pain was well controlled. All patients attended a follow-up visit with the operating surgeon at 4–6 weeks postoperatively.

Results

The mean blood loss was less than 10 cc in both cases and the average operative time was 57 minutes. Both patients showed significant improvement in diarrhea frequency immediately following surgery. No complications or postoperative adverse events were documented at the time of last follow-up.

Weight loss analysis

See Table 1 for weight loss outcomes with SIPS and revision CCL.

Discussion

Being a new procedure, there are very little data describing the rare complications of this unique approach or how to fix them [8,9].

Download English Version:

<https://daneshyari.com/en/article/5661978>

Download Persian Version:

<https://daneshyari.com/article/5661978>

[Daneshyari.com](https://daneshyari.com)