Accepted Manuscript

Do Serial Measurements of Cervical Length Improve the Prediction of Preterm Birth in Asymptomatic Women with Twin Gestations?

Nir Melamed, MD, MSc, Alex Pittini, Liran Hiersch, MD, Yariv Yogev, MD, Steven J. Korzeniewski, Ph.D., Roberto Romero, MD, D.Med.Sci., Jon Barrett, MD

PII: S0002-9378(16)30379-9

DOI: 10.1016/j.ajog.2016.06.034

Reference: YMOB 11171

To appear in: *American Journal of Obstetrics and Gynecology*

Received Date: 9 May 2016

Revised Date: 14 June 2016

Accepted Date: 19 June 2016

Please cite this article as: Melamed N, Pittini A, Hiersch L, Yogev Y, Korzeniewski SJ, Romero R, Barrett J, Do Serial Measurements of Cervical Length Improve the Prediction of Preterm Birth in Asymptomatic Women with Twin Gestations?, *American Journal of Obstetrics and Gynecology* (2016), doi: 10.1016/j.ajog.2016.06.034.

Do Serial Measurements of Cervical Length Improve the Prediction of Preterm Birth in Asymptomatic Women with Twin Gestations?

Nir Melamed, MD, MSc [1], Alex Pittini [1], Liran Hiersch, MD [2], Yariv Yogev, MD [2], Steven J. Korzeniewski Ph.D.[3,4,5], Roberto Romero, MD, D.Med.Sci. [3,5,6,7], Jon Barrett, MD [1]

[1] *Division of Maternal-Fetal Medicine, Department of Obstetrics and Gynecology, Sunnybrook Health Sciences Centre, University of Toronto, Ontario, Canada*
[2] *Department of Obstetrics and Gynecology, Rabin Medical Center, Petach Tikva ,and Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel*
[3] *Perinatology Research Branch, Program for Perinatal Research and Obstetrics, Division of Intramural Research,* Eunice Kennedy Shriver *National Institute of Child Health and Human Development, NIH, Bethesda, MD and Detroit, MI*
[4]*Department of Obstetrics and Gynecology, Wayne State University School of Medicine, Detroit, Michigan*
[5]*Department of Epidemiology and Biostatistics, Michigan State University, East Lansing, MI*
[6]*Department of Obstetrics and Gynecology, University of Michigan, Ann Arbor, MI*
[7]*Center for Molecular Medicine and Genetics, Wayne State University, Detroit, MI*

Correspondence address:

Nir Melamed, MD, MSc

Division of Maternal Fetal Medicine,

Department of Obstetrics and Gynecology ,

Sunnybrook Health Sciences Center, Toronto, ON, Canada

E-mail: nir.melamed@sunnybrook.ca

Disclosure statement: The authors report no conflict of interest.

Sources of financial support: This research was supported, in part, by the Perinatology Research Branch, Division of Intramural Research, *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National