

# Accepted Manuscript

Gestational hypertension is associated with increased risk of type 2 diabetes in adult offspring: The Helsinki Birth Cohort Study

Eero Kajantie, DrMedSc, Clive Osmond, PhD, Johan G. Eriksson, DrMedSc



PII: S0002-9378(16)30977-2

DOI: [10.1016/j.ajog.2016.10.041](https://doi.org/10.1016/j.ajog.2016.10.041)

Reference: YMOB 11371

To appear in: *American Journal of Obstetrics and Gynecology*

Received Date: 9 August 2016

Revised Date: 22 October 2016

Accepted Date: 28 October 2016

Please cite this article as: Kajantie E, Osmond C, Eriksson JG, Gestational hypertension is associated with increased risk of type 2 diabetes in adult offspring: The Helsinki Birth Cohort Study, *American Journal of Obstetrics and Gynecology* (2016), doi: 10.1016/j.ajog.2016.10.041.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Gestational hypertension is associated with increased risk of type 2 diabetes in adult offspring:****The Helsinki Birth Cohort Study**

Eero KAJANTIE, DrMedSc, Clive OSMOND, PhD, Johan G ERIKSSON, DrMedSc

From the Chronic Disease Prevention Unit, National Institute for Health and Welfare, Helsinki, Finland (Drs Kajantie and Eriksson), Hospital for Children and Adolescents, Helsinki University Central Hospital and University of Helsinki, Helsinki, Finland (Dr Kajantie), PEDEGO Research Unit, MRC Oulu, Oulu University Hospital and University of Oulu, Oulu, Finland (Dr Kajantie), MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, United Kingdom (Dr Osmond), Department of General Practice and Primary Health Care, University of Helsinki and Helsinki University Hospital, Helsinki, Finland (Dr Eriksson) and Folkhälsan Research Center, Helsinki, Finland (Dr Eriksson).

The authors report no conflict of interest.

Sources of funding: The study was supported by grants from Academy of Finland [grants 129369, 129907, 135072, 129255 and 126775 to JGE; 127437, 129306, 130326, 134791 and 263924 to EK], Emil Aaltonen Foundation, Finnish Foundation for Cardiovascular Research, Finnish Foundation for Pediatric Research, Juho Vainio Foundation, Finska Läkaresällskapet, Novo Nordisk Foundation, Samfundet Folkhälsan, Liv och Hälsa, Signe and Ane Gyllenberg Foundation, Sigrid Jusélius Foundation, and Yrjö Jahnsson Foundation. None of the funding sources had any role in study design, in the collection, analysis and interpretation of data, in the writing of the report, or in the decision to submit the article for publication.

Corresponding author:

Eero Kajantie

National Institute for Health and Welfare, Chronic Disease Prevention Unit

PL 30, FIN-00271 Helsinki

Finland

Tel. +358-29-5248610

E-mail eero.kajantie@helsinki.fi

Word count: 356 (abstract); 1559 (body text). Figure 1 is to appear in the print publication.

Download English Version:

<https://daneshyari.com/en/article/5676249>

Download Persian Version:

<https://daneshyari.com/article/5676249>

[Daneshyari.com](https://daneshyari.com)