

Accepted Manuscript



Twenty-five percent of patients with clinical chorioamnionitis in preterm gestations have no evidence of either culture-proven intra-amniotic infection or intra-amniotic inflammation

Kyung Joon Oh, MD, PhD, Sun Min Kim, MD, PhD, Joon-Seok Hong, MD, PhD, Eli Maymon, MD, Offer Erez, MD, Bogdan Panaitescu, MD, PhD, Nardhy Gomez-Lopez, PhD, Roberto Romero, MD, D.Med.Sci., Bo Hyun Yoon, MD, PhD

PII: S0002-9378(17)30361-7

DOI: [10.1016/j.ajog.2017.02.035](https://doi.org/10.1016/j.ajog.2017.02.035)

Reference: YMOB 11555

To appear in: *American Journal of Obstetrics and Gynecology*

Received Date: 29 December 2016

Revised Date: 15 February 2017

Accepted Date: 21 February 2017

Please cite this article as: Oh KJ, Kim SM, Hong J-S, Maymon E, Erez O, Panaitescu B, Gomez-Lopez N, Romero R, Yoon BH, Twenty-five percent of patients with clinical chorioamnionitis in preterm gestations have no evidence of either culture-proven intra-amniotic infection or intra-amniotic inflammation, *American Journal of Obstetrics and Gynecology* (2017), doi: 10.1016/j.ajog.2017.02.035.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Twenty-five percent of patients with clinical chorioamnionitis in preterm gestations have no evidence of either culture-proven intra-amniotic infection or intra-amniotic inflammation

Kyung Joon OH, MD, PhD^{1,2}; Sun Min KIM, MD, PhD^{1,3}; Joon-Seok HONG MD, PhD^{1,2}; Eli MAYMON, MD^{4,5}, Offer EREZ, MD^{4,5}, Bogdan PANAITESCU, MD, PhD^{4,5}, Nardhy GOMEZ-LOPEZ, PhD^{4,5,6}, Roberto ROMERO, MD, D.Med.Sci.^{4,7,8,9}; Bo Hyun YOON MD, PhD¹

¹Department of Obstetrics and Gynecology, Seoul National University College of Medicine, Seoul, Republic of Korea; ²Department of Obstetrics and Gynecology, Seoul National University Bundang Hospital, Seongnam-si, Republic of Korea; ³Department of Obstetrics and Gynecology, Seoul Metropolitan Government Seoul National University Boramae Medical Center, Seoul, Republic of Korea; ⁴Perinatology Research Branch, NICHD/NIH/DHHS, Bethesda, Maryland, USA, and Detroit, Michigan, USA; ⁵Department of Obstetrics and Gynecology, Wayne State University School of Medicine, Detroit, Michigan, USA; ⁶Department of Immunology and Microbiology, Wayne State University School of Medicine, Detroit, MI, USA; ⁷Department of Obstetrics and Gynecology, University of Michigan, Ann Arbor, Michigan, USA; ⁸Department of Epidemiology and Biostatistics, Michigan State University, East Lansing, Michigan, USA; ⁹Center for Molecular Medicine and Genetics, Wayne State University, Detroit, Michigan, USA.

Conflict of Interest: The authors report no conflicts of interest.

Source of Funding: This work was supported in part, by the SNUH research fund (Grant No. 04-2016-0920), and in part, by the Perinatology Research Branch, Program for Perinatal Research and Obstetrics, Division of Intramural Research, *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National Institutes of Health, U.S. Department of Health and Human Services (NICHD/NIH/DHHS), and, in part, with Federal funds from NICHD/NIH/DHHS under Contract No. HHSN275201300006C.

Download English Version:

<https://daneshyari.com/en/article/5676310>

Download Persian Version:

<https://daneshyari.com/article/5676310>

[Daneshyari.com](https://daneshyari.com)