

# Accepted Manuscript

While mortality rates differ following dysvascular partial foot and transtibial amputation, should they influence the choice of amputation level?

Michael Dillon, PhD, Stefania Fatone, PhD, Matthew Quigley, MCPO(Hons)

PII: S0003-9993(17)30254-X

DOI: [10.1016/j.apmr.2017.04.002](https://doi.org/10.1016/j.apmr.2017.04.002)

Reference: YAPMR 56864

To appear in: *ARCHIVES OF PHYSICAL MEDICINE AND REHABILITATION*

Received Date: 25 February 2017


Revised Date: 22 March 2017

Accepted Date: 8 April 2017

Please cite this article as: Dillon M, Fatone S, Quigley M, While mortality rates differ following dysvascular partial foot and transtibial amputation, should they influence the choice of amputation level?, *ARCHIVES OF PHYSICAL MEDICINE AND REHABILITATION* (2017), doi: 10.1016/j.apmr.2017.04.002.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.





Mortality. It's due to older age  
and advanced systemic disease,  
**NOT** the choice of amputation level.

Download English Version:

<https://daneshyari.com/en/article/5677422>

Download Persian Version:

<https://daneshyari.com/article/5677422>

[Daneshyari.com](https://daneshyari.com)